

Swarthmore College

Works

Physics & Astronomy Faculty Works

Physics & Astronomy

7-20-2003

X-Ray Emission-Line Profile Modeling Of O Stars: Fitting A Spherically Symmetric Analytic Wind-Shock Model To The Chandra Spectrum Of Zeta Puppis

Roban Hultman Kramer , '03

David H. Cohen

Swarthmore College, dcohen1@swarthmore.edu

S. P. Owocki

Follow this and additional works at: <https://works.swarthmore.edu/fac-physics>



Part of the [Astrophysics and Astronomy Commons](#)

Let us know how access to these works benefits you

Recommended Citation

Kramer, Roban Hultman , '03; Cohen, David H.; and Owocki, S. P.. Roban Hultman Kramer , '03 et al. (2003). "X-Ray Emission-Line Profile Modeling Of O Stars: Fitting A Spherically Symmetric Analytic Wind-Shock Model To The Chandra Spectrum Of Zeta Puppis". *Astrophysical Journal*. Volume 592, Issue 1. 532-538. DOI: 10.1086/375390

<https://works.swarthmore.edu/fac-physics/13>

This work is brought to you for free by Swarthmore College Libraries' Works. It has been accepted for inclusion in Physics & Astronomy Faculty Works by an authorized administrator of Works. For more information, please contact myworks@swarthmore.edu.

X-RAY EMISSION-LINE PROFILE MODELING OF O STARS: FITTING A SPHERICALLY SYMMETRIC ANALYTIC WIND-SHOCK MODEL TO THE *CHANDRA* SPECTRUM OF ζ PUPPIS

ROBAN H. KRAMER,^{1,2} DAVID H. COHEN,¹ AND STANLEY P. OWOCKI³

Received 2002 November 22; accepted 2003 March 17

ABSTRACT

X-ray emission-line profiles provide the most direct insight into the dynamics and spatial distribution of the hot, X-ray-emitting plasma above the surfaces of OB stars. The O supergiant ζ Puppis shows broad, blueshifted, and asymmetric line profiles, generally consistent with the wind-shock picture of OB star X-ray production. We model the profiles of eight lines in the *Chandra* HETGS spectrum of this prototypical hot star. The fitted lines indicate that the plasma is distributed throughout the wind starting close to the photosphere, that there is significantly less attenuation of the X-rays by the overlying wind than is generally supposed, and that there is not a strong trend in wind absorption with wavelength.

Subject headings: line: profiles — stars: early-type — stars: individual (ζ Puppis) — stars: mass loss — stars: winds, outflows — X-rays: stars

1. INTRODUCTION

The nature of the copious soft X-ray emission from hot stars has been a long-standing controversy since its discovery in the late 1970s (Harnden et al. 1979; Cassinelli & Olson 1979). Solar-type coronal emission was first assumed (Cassinelli & Olson 1979; Waldron 1984), but wind-shock models of various types gained currency throughout the following decade (Waldron 1984; Owocki, Castor, & Rybicki 1988; MacFarlane & Cassinelli 1989; Chen & White 1991; Hillier et al. 1993; Cohen et al. 1996; Feldmeier et al. 1997a; Feldmeier, Puls, & Pauldrach 1997b). More recently, hybrid magnetic wind models have been proposed for some hot stars (Gagné et al. 1997; Babel & Montmerle 1997; ud-Doula & Owocki 2002).

Until the launch of *Chandra* and *XMM-Newton*, no observational diagnostics were available that could provide a direct discriminant between the coronal and wind-shock paradigms. However, because of their superior spectral resolution, both of these new telescopes allow for the separation of individual emission lines and the resolution of Doppler-broadened line profiles. The spectral resolution of *Chandra*'s grating spectrometers exceeds $\lambda/\Delta\lambda \sim 1000$ (for the FWHM), which corresponds to a velocity of 300 km s^{-1} , and that of the *XMM-Newton* RGS is almost as great. This compares favorably to the terminal velocities of the radiation-driven winds of O stars, which approach $v_\infty = 3000 \text{ km s}^{-1}$, implying ~ 20 resolution elements for a velocity range of $2v_\infty$.

At the most basic level, the X-ray emission lines from hot stars will be either narrow and therefore roughly consistent with coronal emission or broad and roughly consistent with wind-shock emission. This is because in the wind-shock model the high velocities of X-ray-emitting plasma embedded in the wind would Doppler shift the emission

across a range of wavelengths. Initial papers reporting on *Chandra* and *XMM-Newton* observations of various O and B stars (Schulz et al. 2001; Kahn et al. 2001; Waldron & Cassinelli 2001; Cassinelli et al. 2001; Miller et al. 2002; Cohen et al. 2003) discussed line widths, which vary from large for early O stars to small for B stars. Some of these initial studies noted that the emission lines can be blueshifted and the profiles somewhat asymmetric, but none of these studies discussed or modeled the shapes of the resolved emission lines.

In this paper we fit a specific model of X-ray emission-line profiles in an expanding, emitting, and absorbing wind to a *Chandra* HETGS/MEG spectrum of ζ Puppis. The model we fit is empirical and flexible, with only three free parameters. The model assumes a two-component fluid, having as its major constituent the cold, X-ray-absorbing plasma that gives rise to the characteristic UV absorption lines observed in hot star winds, and as its minor constituent the hot, X-ray-emitting plasma. This empirical model is not tied to any one specific physical model of X-ray production and is general enough to fit data representative of any of the major models, so long as they are spherically symmetric. To the extent that wind-shock models are found to be consistent with the observed line profiles, our model parameters can be used to constrain the physical properties of the shock-heated plasma. This ultimately can be used to constrain the values of the physical parameters of the appropriate wind-shock model.

In § 2 we discuss the physical effects leading to nontrivial line shapes and describe the specific empirical model we use to perform the fits. In § 3 we describe the *Chandra* ζ Puppis data set and how we perform the line fitting and parameter estimation, and in § 4 we discuss the derived model parameters in the context of the various physical models that have been proposed to explain hot star X-ray emission, as well as in the context of other X-ray diagnostics that have recently been applied to the data from this star.

2. THEORETICAL CONSIDERATIONS

In the context of the fast, radiation-driven winds of OB stars, a source of X-ray emission embedded in the wind will

¹ Swarthmore College Department of Physics and Astronomy, 500 College Avenue, Swarthmore, PA 19081; roban@scs.swarthmore.edu, cohen@hven.swarthmore.edu.

² Prism Computational Sciences, 455 Science Drive, Madison, WI 53711.

³ Bartol Research Institute, University of Delaware, 217 Sharp Laboratory, Newark, DE 19716; owocki@bartol.udel.edu.

lead to Doppler-broadened profiles, but only the X-ray-emitting plasma traveling at the wind terminal velocity directly toward or away from the observer will lead to maximal blueshifts and redshifts. The amount of emission at each intermediate wavelength, and thus the shape and characteristic width of the line, depends on the spatial and velocity distribution of the hot plasma. As described by MacFarlane et al. (1991) in the case of a shell of X-ray-emitting plasma in a hot star wind, the continuum absorption of X-rays by the cool component of the wind will cause the resulting emission lines to be attenuated on the red side and be relatively unaffected on the blue side of the line profile. The apparent peak of the emission line thus shifts to the blue side of the line center, and the line is asymmetric, with a shallower red wing and a steeper blue wing.

This basic idea of Doppler-broadened emission from a hot wind component and continuum absorption by the cool wind component leading to a broadened, shifted, and asymmetric line was extended from a shell to a spherically symmetric wind by Ignace (2001). He showed that model line profiles could be generated analytically for a constant-velocity wind. Owocki & Cohen (2001) extended this concept further, to an accelerating wind, with a model having four free parameters. Two describe the spatial distribution of the X-ray-emitting plasma: R_0 , the minimum radius of X-ray emission, and q , the radial power-law index of the emissivity. There is assumed to be no emission below $r = R_0$. Above $r = R_0$, volume emissivity is assumed to scale like the density of the wind squared ρ^2 (since collisional processes and recombination dominate the ionization/excitation kinematics), with an extra factor r^{-q} allowing for spatial variation of shock temperatures, cooling structures, and density and filling factor of the shocked material. The parameter β controls the velocity of the wind, which is assumed to follow a “beta-velocity law”:

$$v(r) = v_\infty(1 - R_*/r)^\beta. \quad (1)$$

Both components of the wind follow the same velocity law in all cases discussed here, but in principle they could be allowed to differ. The fourth parameter, τ_* , characterizes the amount of absorption in the wind,

$$\tau_* \equiv \frac{\kappa \dot{M}}{4\pi v_\infty R_*}, \quad (2)$$

where κ is the line opacity, or mass absorption coefficient ($\text{cm}^2 \text{s}^{-1}$). This relates to the commonly quoted radius of optical depth unity R_1 by the equation (for $\beta = 1$)

$$\frac{R_1}{R_*} = \frac{1}{1 - \exp(-1/\tau_*)} \approx \tau_* + 0.5 \quad \text{for } \tau_* > 0.5. \quad (3)$$

The error in the above approximation is less than 10% for $\tau_* > 0.67$ (or $R_1/R_* > 1.3$).

The model put forward by Owocki & Cohen (2001) is a phenomenological one. It describes the physical properties of the hot and cool components of the wind, but does not describe the physics underlying the generation of the hot plasma. It is therefore quite flexible, capable of describing a thin shell of X-ray-emitting plasma, including a coronal-type zone near the photosphere, as well as wind shocks distributed spatially throughout the wind, with the shock distribution and wind velocity varying with radius. This model is thus capable of constraining properties of both the

X-ray-emitting and the X-ray-absorbing wind components. When applied to an ensemble of lines it has the potential to constrain these properties as a function of both temperature and wavelength. Assuming turbulent and thermal broadening are negligible and combining the other model assumptions, the line profile as a function of scaled wavelength $x \equiv (\lambda/\lambda_0 - 1)(c/v_\infty)$ is given by

$$L_x \propto \int_{r=r_x}^{\infty} \frac{r^{-(q+2)}}{(1 - R_*/r)^{3\beta}} \exp[-\tau(\mu_x, r)] dr, \quad (4)$$

where $r_x \equiv \max[R_0, R_*/(1 - |x|^{1/\beta})]$, $\mu_x \equiv x/(1 - R_*/r)^\beta$, and $\tau(\mu, r)$ (which is proportional to τ_*) is the optical depth along the observer’s line of sight at direction $\cos \mu$ and radial coordinate r . The constant of proportionality (which is itself proportional to the emission measure) will not be determined in this work. Equation (4) must be solved numerically, except in the case of $\beta = 0$ (constant-velocity wind). Even so, we only obtain solutions for integer values of β .

This model assumes implicitly that the sites of X-ray emission are so numerous and well mixed with the primary cool wind component that we can treat the wind as a two-component fluid. It also neglects nonradial velocity components, including small-scale fluctuations like turbulence. Note that the assumption of purely radial velocities is what allows wind absorption to break the symmetry of the line, since redshifted emission always arises at higher line-of-sight distances (and therefore higher optical depths) than the blueshifted emission (see the contour plots in Figs. 1 and 2).

Similar treatments of radiation transport have previously been fitted to the global form of low-resolution spectra (Hillier et al. 1993; Feldmeier et al. 1997a). MacFarlane et al. (1991) and Waldron & Cassinelli (2001) have modeled line profiles from discrete spherical shocks. Elsewhere we report on the investigation of nonspherical models and their applicability to hot stars (Kramer et al. 2003; Tonnesen et al. 2003).

For this study, we have adopted the Owocki & Cohen (2001) model and performed fits on eight strong lines in the *Chandra* HETGS spectrum of the O4 supergiant ζ Puppis, extracting best-fit values and associated confidence limits for three model parameters: q , R_0 , and τ_* .

3. FITTING THE MODEL TO OBSERVED LINE PROFILES

Our data set consists of the ± 1 order MEG spectrum from a 67 ks observation of the O4f star ζ Puppis first reported on by Cassinelli et al. (2001). The FWHM of the MEG spectral response is $\Delta\lambda_{\text{MEG}} = 0.023 \text{ \AA}$ (*Chandra* X-Ray Center 2001).⁴ All the distinguishable lines in this spectrum are many times more broad, allowing their profiles to be well resolved. The breadth of the lines means that many of them are contaminated by emission from neighboring lines. After eliminating He-like forbidden, intercombination, and resonance triplets as unsuitable for fitting due to excessive blending, we identified other potential blends by visual inspection of the spectrum and by referring to the line strengths calculated by Mewe, Gronenschild, & van den

⁴ The *Chandra* Proposers’ Observatory Guide is available at <http://cxc.harvard.edu/udocs/docs/>.




FIG. 1.—Two representative lines with best-fit models. Shown are *Chandra* MEG spectra of (a) Fe xvii at 15.01 Å and (c) O viii at 18.97 Å (in gray), with corresponding best-fit profiles (in black). Laboratory rest wavelengths are indicated by the vertical dashed lines. To the right we show (b and d) contour plots representing the models. The observer is located at ($p/R_* = 0, z/R_* = \infty$). The inner circle is of radius R_* , the outer circle of radius R_0 . Gray contours are curves of constant line-of-sight velocity component in units of v_∞ . Dashed contours are curves of constant optical depth ($\tau = 0.5, 1, 2$), integrated along the line of sight.

Oord (1985, Table 4) and those in the Astrophysical Plasma Emission Database (APED; Smith et al. 2001).⁵ We consider a line with rest wavelength λ_0 to extend over a wavelength range defined by

$$\lambda_0 \left(1 + \frac{v_\infty}{c}\right) + \Delta\lambda_{\text{MEG}} \geq \lambda \geq \lambda_0 \left(1 - \frac{v_\infty}{c}\right) - \Delta\lambda_{\text{MEG}}. \quad (5)$$

⁵ The Interactive GUIDE for ATOMDB is available at <http://obsvis.harvard.edu/WebGUIDE/>.

The widths of neighboring lines are calculated the same way, and any overlap in the ranges is excluded from the fit (see Table 1 for the wavelength range over which each fit was performed). We adopted the terminal velocity value determined by Prinja, Barlow, & Howarth (1990), $v_\infty = 2485 \text{ km s}^{-1}$.

We numerically integrate equation (4) over the desired wavelength range using software written in Mathematica. The resulting profile is convolved with a Gaussian




FIG. 2.—Models at extremes of the confidence region. Shown are the *Chandra* MEG spectrum of the 17.054 Å line of Fe xvii (in gray), and (in black) the best-fit model ($\tau_* = 0.5$), the fit with τ_* held at its 95.4% confidence upper limit ($\tau_* = 1.5$) and the fit with τ_* held at its 95.4% confidence lower limit ($\tau_* = 0.0$). The contour plots are the same style as in Fig. 1 and correspond to (b) the $\tau_* = 0.0$ model, (c) the best-fit model, and (d) the $\tau_* = 1.5$ model.

representing instrumental response, binned identically to the data, and normalized to predict the same total number of counts as were observed over the same wavelength range.

To properly treat the statistics of Poisson-distributed, low photon count data, we use Cash's C (Cash 1979) as the fit statistic. Fits are performed by calculating C on a grid in parameter space. The coordinates of the parameter space point producing the minimum C value C_{\min} are taken as our best-fit parameters. We then use limits on $\Delta C = C - C_{\min}$ to define our confidence regions, as described by Cash (1979). The grid is expanded as needed until the entire confidence region is encompassed within it. In all our fits we held

β constant at $\beta = 1$ (which is very close to the typical O-star value of 0.8; Groenewegen, Lamers, & Pauldrach 1989) and varied R_0 , q , and τ_* .

To confirm the confidence limits derived using the ΔC statistic, we carried out the fitting procedure on Monte Carlo simulated data sets and compared the parameter-space distribution of simulated-data fit parameters to the calculated ΔC confidence region for each line. The Monte Carlo simulations gave results that were consistent with those given by the ΔC statistic.

The quality of the fits was evaluated using the Kuiper statistic, a variant of the Kolmogorov-Smirnov test.

TABLE 1
BEST-FIT PARAMETERS WITH 95.4% CONFIDENCE LIMITS

Ion	λ_0 (Å)	$\lambda_0/\Delta\lambda$	$2v_\infty/\Delta v$	q	R_0	τ_*	x_{\min}	x_{\max}	N_{obs}	α_0	α_{RF}
N VII.....	24.78	1077	18	$-0.5^{+0.6}_{-0.3}$	$2.0^{+1.3}_{-0.6}$	$0.5^{+2.0}_{-0.5}$	-1.11	1.11	92	0.667	0.356
O VIII.....	18.97	825	14	$-0.1^{+0.6}_{-0.4}$	$1.2^{+2.1}_{-0.2}$	$2.5^{+2.5}_{-1.5}$	-1.04	1.14	69	0.941	0.790
Fe XVII.....	17.054	741	12	$-0.6^{+0.4}_{-0.2}$	$1.4^{+0.6}_{-0.3}$	$0.5^{+1.0}_{-0.5}$	-0.74	1.16	54	0.971	0.950
Fe XVII ^a	16.787	730	12	$0.4^{+0.2}_{-0.6}$	$1.0^{+2.3}_{-0.0}$	$4.5^{+3.5}_{-2.5}$	-1.16	0.74	53	0.517	0.490
Fe XVII.....	15.262	664	11	$-0.8^{+0.2}_{-0.2}$	$1.4^{+1.1}_{-0.4}$	$1.5^{+2.5}_{-1.5}$	-0.81	1.19	50	0.039	0.151
Fe XVII.....	15.013	653	11	$-0.2^{+0.4}_{-0.3}$	$1.4^{+0.6}_{-0.3}$	$1.0^{+1.0}_{-0.5}$	-1.19	0.79	49	0.887	0.800
Ne X.....	12.132	627	9	$-0.2^{+0.8}_{-0.8}$	$1.3^{+0.7}_{-0.3}$	$2.4^{+1.2}_{-1.8}$	-1.23	0.08	26	0.609	0.370
Si XIV ^b	6.18	269	4	$-0.2^{+0.0}_{-0.8}$	$1.4^{+8.6}_{-1.4}$	$1.5^{+5.5}_{-1.5}$	-0.98	0.59	16	0.857	0.622

NOTES.—The width of the instrumental response in wavelength units is $\Delta\lambda = \Delta\lambda_{\text{MEG}} = 0.023 \text{ \AA}$, or in velocity units $\Delta v = c\Delta\lambda/\lambda_0$. The scaled wavelength $x \equiv (c/v_\infty)(\lambda - \lambda_0)/\lambda_0$. N_{obs} is the number of wavelength bins included in the fit.

^a The anomalous results for these lines may be due to contamination.

^b At the 95.4% confidence level upper limit, q is unconstrained for this fit.

Significance levels were determined using Monte Carlo techniques. We obtained two slightly different distributions of the Kuiper statistic depending on whether the simulated data sets were compared directly to the parent model (giving significance level α_0) or to a new best-fit model found by performing the fitting procedure on the simulated data set (giving significance level α_{RF}).

In all, we fit eight lines between 6.18 Å (Si XIV) and 24.78 Å (N VII). The results are listed in Table 1, and shown in Figure 1 for two representative lines. All the fits but one are formally good according to both distributions. The value of $\alpha_0 = 0.039$ for the 15.262 Å fit does not meet the criterion $\alpha \geq 0.05$ for a formally good fit. This line may be contaminated by the Fe XIX lines at 15.198 and 15.3654 Å (APED). The relatively low significance values for the N VII line may simply be the result of random variation or could be a sign that there are resolved spectral features not explained by this simple model. In any case, the fit is formally good. The 16.787 Å fit may also be affected by a blend with an Fe XIX line, this one at 16.718 Å. This would add flux to the blue edge of the line, increasing its skew and explaining the relatively high values of τ_* and q . The Fe XVII line at 17.054 Å is likely blended with another very near by Fe XVII line at 17.096 Å, which may have a small effect on the overall width of the feature.

To demonstrate the typical range of models that can be fitted to one line, we show the best-fit and two extreme models superposed on the Fe XVII line at 17.05 Å in Figure 2. The two extreme models are for the parameter sets that have the largest and smallest values of τ_* within the 95.4% confidence region.

Finally, we summarize the best fit and 95.4% confidence limits of the three free model parameters for seven of the eight lines in Figure 3 (the values for the Si XIV line are not shown because it is at a much shorter wavelength and its fit parameters are poorly constrained). Trends in these fitted parameters and their implications are discussed in the next section.

4. DISCUSSION

The primary result of the analysis presented in this paper is that the X-ray emission lines in the prototypical O supergiant, ζ Puppis, can, for the most part, be adequately fitted with a spherically symmetric wind model having a small number of free parameters. Furthermore, the derived

parameters are quite reasonable in the context of most wind-shock models, being consistent with hot plasma uniformly distributed throughout the wind above a moderate onset radius, X-ray-emitting plasma extending out to the wind terminal velocity, and the need for the inclusion of some wind attenuation.

In detail, however, some interesting trends emerge. First of all, the amount of wind attenuation is significantly smaller than what one might expect from a spherically symmetric smooth wind, given what is known about this star's mass-loss rate and wind opacity. There have been various calculations of the wind optical depth (often expressed as the radius of optical depth unity) as a function of wavelength for this star. They range from values much larger than what we derive here ($7 < \tau_* < 30$ calculated by Hillier et al. 1993, using $\dot{M} = 5.0 \times 10^{-6} M_\odot \text{ yr}^{-1}$, $R_* = 19 R_\odot$, $v_\infty = 2200 \text{ km s}^{-1}$) to values modestly larger ($4 < \tau_* < 8$ calculated by Cassinelli et al. 2001, using values from Lamers & Leitherer 1993: $\dot{M} = 2.4 \times 10^{-6} M_\odot \text{ yr}^{-1}$, $R_* = 16 R_\odot$, $v_\infty = 2200 \text{ km s}^{-1}$). Note that Hillier et al. (1993) find different values for R_1 depending on whether helium recombines or remains ionized in the outer wind, but at energies above 0.5 keV (where all of the lines presented here occur) there is little difference between the two scenarios. More recent stellar parameters determined by Puls et al. (1996) ($\dot{M} = 5.9 \times 10^{-6} M_\odot \text{ yr}^{-1}$, $R_* = 19 R_\odot$, and $v_\infty = 2250 \text{ km s}^{-1}$) agree well with the values used by Hillier et al. (1993) but would increase the Cassinelli et al. (2001) τ_* values by a factor of 2, given the same opacity (see eq. [2]).

If we accept the τ_* values derived from our fits, then the disparity between those values and the ones mentioned above suggests that either the mass-loss rates or wind opacities are being overestimated in previous calculations. The mass-loss rate of ζ Puppis is by now quite well established using UV absorption lines and H α , although improper ionization corrections or clumping could lead to systematic errors. The wind opacity determination seems much more uncertain, both because of the inconsistent values in the literature and because of the difficulty in determining the ionization state of the wind (MacFarlane et al. 1993; MacFarlane, Cohen, & Wang 1994). Recent advances in stellar atmosphere modeling may help to improve these determinations (Pauldrach, Hoffmann, & Lennon 2001). Another means of lowering the wind attenuation is to clump the wind into small clouds that are individually optically thick, rendering the wind porous and enhancing the escape




FIG. 3.—Best-fit values of (a) q , (b) R_0 in units of R_* , and (c) τ_* for seven of the eight lines (filled circles), along with the range given by the 95.4% confidence limits (triangles). See Table 1.

probability of X-ray photons, thus lowering the mean wind opacity. This would also affect the mass-loss rate diagnostics, but is itself an independent effect.

An even more curious result of the τ_* fits is that they are nearly independent of wavelength. This is surprising, because photoionization cross sections should scale roughly as a power of wavelength between λ^2 and λ^3 (Hillier et al. 1993). It is possible that the distribution of ionization edges could conspire to make this relationship much flatter over a small range of wavelengths (as the calculations from Cassinelli et al. 2001 seem to indicate). However, wind clumping might play some role here, too. If the wind opacity is dominated by clumps that are individually optically thick across the wavelength range, then the opacity ceases to be a function of wavelength and instead depends on the physical cross sections of the clumps themselves. We note that the UV line opacity necessary to explain the observed absorption line profiles could, in principle, still be provided by the

tenuous interclump wind, as the line cross sections are much larger than the X-ray photoionization cross sections.

If $R_1 \gg R_0$, the line profile is insensitive to changes in R_0 , since emission much below R_1 is largely absorbed by the wind. The values we find for R_0 , although generally small, are comparable to our values of R_1 (from τ_* by eq. [3]). It is hard to assess these relatively small onset radii in the context of the small (sometimes surprisingly small) values claimed on the basis of observed f/i ratios in He-like ions (Cassinelli et al. 2001; Kahn et al. 2001; Waldron & Cassinelli 2001). This is partially because we do not fit the profiles of any He-like lines (they are too blended) and partly because the most extreme results (smallest value for R_1) are for S xv, which is a higher ionization stage than any of the lines we fit.

The fit results for the parameter q indicate that there is not a strong radial trend in the filling factor. One might expect some competition in a wind shock model between the

tendency to have more and stronger shocks near the star, where the wind is still accelerating, and the tendency for shock heated gas to cool less efficiently in the far wind, where densities are low. Perhaps these two effects cancel to give the observed $q \approx 0$ relationship.

In conclusion, the simple, spherically symmetric wind shock model is remarkably consistent with the observed line profiles in ζ Puppis, providing the most direct evidence yet that some type of wind-shock model applies to this hot star. However, there are indications that the absorption properties of the wind of ζ Puppis, and perhaps other hot stars, must be reconsidered. We will fit this same model to the *Chandra* spectra of other hot stars in the future. However, the lack of strong line asymmetries in stars such as ζ Ori and δ Ori and the narrow lines in θ^1 Ori C and τ Sco indicate that spheri-

cally symmetric wind-shock models with absorption may not fit the data from these stars as well as they do ζ Puppis.

We wish to acknowledge grants GO0-1089A and G02-3024B to Swarthmore College and Prism Computational Sciences. R. H. K. acknowledges funding provided by a Howard Hughes Medical Institute undergraduate research grant. S. P. O. acknowledges partial support by NSF grant AST 00-97983 and NASA grant NAG 5-3530 to the Bartol Research Institute of the University of Delaware. We would also like to thank the referee, Rolf-Peter Kudritzki, for his comments, which substantially improved the paper, and Elliot Reed of Swarthmore College for his initial important work on this project.

REFERENCES

- Babel, J., & Montmerle, T. 1997, *ApJ*, 485, L29
 Cash, W. 1979, *ApJ*, 228, 939
 Cassinelli, J. P., Miller, N. A., Waldron, W. L., MacFarlane, J. J., & Cohen, D. H. 2001, *ApJ*, 554, L55
 Cassinelli, J. P., & Olson, G. L. 1979, *ApJ*, 229, 304
Chandra X-Ray Center 2001, *The Chandra Proposers' Observatory Guide*, version 4.0 (Cambridge: *Chandra* X-Ray Center)
 Chen, W., & White, R. L. 1991, *ApJ*, 366, 512
 Cohen, D. H., Cooper, R. G., MacFarlane, J. J., Owocki, S. P., Cassinelli, J. P., & Wang, P. 1996, *ApJ*, 460, 506
 Cohen, D. H., De Mèssieres, G. E., MacFarlane, J. J., Miller, N. A., Cassinelli, J. P., Owocki, S. P., & Liedahl, D. A. 2003, *ApJ*, 586, 495
 Feldmeier, A., Kudritzki, R.-P., Palsa, R., Pauldrach, A. W. A., & Puls, J. 1997a, *A&A*, 320, 899
 Feldmeier, A., Puls, J., & Pauldrach, A. W. A. 1997b, *A&A*, 322, 878
 Gagné, M., Caillault, J.-P., Stauffer, J. R., & Linsky, J. L. 1997, *ApJ*, 478, L87
 Groenewegen, M. A. T., Lamers, H. J. G. L. M., & Pauldrach, A. W. A. 1989, *A&A*, 221, 78
 Harnden, F. R., Jr., et al. 1979, *ApJ*, 234, L51
 Hillier, D. J., Kudritzki, R. P., Pauldrach, A. W., Baade, D., Cassinelli, J. P., Puls, J., & Schmitt, J. H. M. M. 1993, *A&A*, 276, 117
 Ignace, R. 2001, *ApJ*, 549, L119
 Kahn, S. M., Leutenegger, M. A., Cotam, J., Rauw, G., Vreux, J.-M., den Boggende, A. J. F., Mewe, R., & Güdel, M. 2001, *A&A*, 365, L312
 Kramer, R. H., Tonnesen, S. K., Cohen, D. H., Owocki, S. P., ud-Doula, A., & MacFarlane, J. J. 2003, *Rev. Sci. Instrum.*, 74, 1966
 Lamers, H. J. G. L. M., & Leitherer, C. 1993, *ApJ*, 412, 771
 MacFarlane, J. J., & Cassinelli, J. P. 1989, *ApJ*, 347, 1090
 MacFarlane, J. J., Cassinelli, J. P., Welsh, B. Y., Vedder, P. W., Vallerger, J. V., & Waldron, W. L. 1991, *ApJ*, 380, 564
 MacFarlane, J. J., Cohen, D. H., & Wang, P. 1994, *ApJ*, 437, 351
 MacFarlane, J. J., Waldron, W. L., Corcoran, M. F., Wolff, M. J., Wang, P., & Cassinelli, J. P. 1993, *ApJ*, 419, 813
 Mewe, R., Gronenschild, E. H. B. M., & van den Oord, G. H. J. 1985, *A&AS*, 62, 197
 Miller, N. A., Cassinelli, J. P., Waldron, W. L., MacFarlane, J. J., & Cohen, D. H. 2002, *ApJ*, 577, 951
 Owocki, S. P., Castor, J. I., & Rybicki, G. B. 1988, *ApJ*, 335, 914
 Owocki, S. P., & Cohen, D. H. 2001, *ApJ*, 559, 1108
 Pauldrach, A. W. A., Hoffmann, T. L., & Lennon, M. 2001, *A&A*, 375, 161
 Prinja, R. K., Barlow, M. J., & Howarth, I. D. 1990, *ApJ*, 361, 607
 Puls, J., et al. 1996, *A&A*, 305, 171
 Schulz, N. S., Canizares, C. R., Huenemoerder, D., & Lee, J. C. 2001, *ApJ*, 549, 441
 Smith, R. K., Brickhouse, N. S., Liedahl, D. A., & Raymond, J. C. 2001, *ApJ*, 556, L91
 Tonnesen, S. K., Cohen, D. H., Owocki, S. P., ud-Doula, A., Gagne, M., & Oksala, M. 2003, *BAAS*, 34, 1284
 ud-Doula, A., & Owocki, S. P. 2002, *ApJ*, 576, 413
 Waldron, W. L. 1984, *ApJ*, 282, 256
 Waldron, W. L., & Cassinelli, J. P. 2001, *ApJ*, 548, L45