

12-1-2015

Using Protoplanetary Disks To Weigh The Youngest Stars And Constrain The Earliest Stages Of Stellar Evolution

I. Czekala

S. Andrews

Eric L. N. Jensen

Swarthmore College, ejensen1@swarthmore.edu

K. Stassun

D. W. Latham

See next page for additional authors

Follow this and additional works at: <http://works.swarthmore.edu/fac-physics>

 Part of the [Astrophysics and Astronomy Commons](#)

Recommended Citation

I. Czekala, S. Andrews, Eric L. N. Jensen, K. Stassun, D. W. Latham, D. Wilner, and G. Torres. (2015). "Using Protoplanetary Disks To Weigh The Youngest Stars And Constrain The Earliest Stages Of Stellar Evolution". *Bulletin Of The American Astronomical Society*.

Volume 47, Issue 6.

<http://works.swarthmore.edu/fac-physics/243>

This Poster Session is brought to you for free and open access by the Physics & Astronomy at Works. It has been accepted for inclusion in Physics & Astronomy Faculty Works by an authorized administrator of Works. For more information, please contact myworks@swarthmore.edu.

Authors

I. Czekala, S. Andrews, Eric L. N. Jensen, K. Stassun, D. W. Latham, D. Wilner, and G. Torres

Extreme Solar Systems III Meeting
Big Island, HI – November/December, 2015
Meeting Abstracts

Session Table of Contents

100 – Overview of Observations	Session	Interactions
101 – Radial Velocities	113 – Planet Detection - Radial Velocities	201 – Dynamical Evolution
102 – Transiting Planets I	Poster Session	202 – Direct Imaging I
103 – Transiting Planets II	114 – Planet Detection - Transits Poster	203 – Direct Imaging II
104 – Direct Imaging Poster Session	Session	300 – Planet Formation
105 – Disks and Migration Poster Session	115 – Planet Formation and Interior	301 – Structure and Evolution
106 – Future Missions and	Structure Modeling Poster Session	400 – Atmospheres I
Instrumentation Poster Session	116 – Planets around Compact Objects	401 – Atmospheres II
107 – Habitability Poster Session	Poster Session	402 – Planets in and around Binaries I
108 – Hot Jupiters and Star-Planet	117 – Planets in and around Binary Stars	403 – Planets in and around Binaries II
Interactions Poster Session	Poster Session	500 – Habitability and Biosignatures
109 – Orbital Dynamics and Planet-Planet	118 – Population Statistics and	501 – Population Statistics and
Interactions Poster Session	Mass-Radius Relations Poster Session	Mass-Radius Relations
110 – Other Topics Poster Session	119 – Super-Earths and Mini-Neptunes	502 – Planets around Evolved Stars and
111 – Planetary Atmospheres - Close-In and	Poster Session	Compact Remnants
Transiting Gas Giants Poster Session	120 – Young Systems Poster Session	503 – TESS and Other Future Missions
112 – Planet Detection - Other Poster	200 – Ultrashort Periods and Planet-Star	

resonances in exoplanet pairs

Mean-motion resonances (MMRs) are typically stable configurations for pairs of planets. Given that planets should migrate relative to one another in their natal disk, one might expect to have found most planets locked in such MMRs. The fact that most Kepler planets are *not* observed in MMRs therefore requires an explanation. Goldreich and Schlichting (2014) recently argued that, in fact, due to interactions with the protoplanetary disk, planets below a threshold mass should break out of the strongest MMRs, i.e., the MMRs become *overstable*.

While follow-up work has studied the robustness of this result to varying orbital architectures, we focus on the specific numerical implementation of the disk effects, which translates into differing physical interpretations of the planet-disk interactions. We will present how these physical choices affect the parameter space in which overstability sets in, and how certain choices can generate spurious results. We will then extend our results to general cases of broad applicability, and summarize the merits and pitfalls of these different numerical implementations of perturbations from the protoplanetary disk, particularly in tightly packed systems.

We have packaged these numerical implementations into REBOUNDx, an open-source C and Python package for incorporating planet-disk interactions, as well as additional effects (like post-newtonian corrections), into N-body simulations using REBOUND. We will give a brief demo that highlights its ease of installation and use, as well as its synergy with Python's powerful plotting and scientific analysis libraries.

Author(s): Daniel Tamayo², Hanno Rein², Alice Chen², morgan bennett¹
Institution(s): 1. *Simon Fraser University*, 2. *University of Toronto at Scarborough*

105.03 – Constraints on Exoplanet System Architectures from Debris Disks

Debris disks are dusty disks around main sequence stars. Terrestrial planets may be forming in young debris disks with ages <100 Myr. Planets in debris disks dynamically sculpt the dust in these systems. Thus, the spatial structure of debris disks could be an indicator of where planets have formed. We present an analysis of several members of the Scorpius-Centaurus OB Association (Sco Cen) that host both debris disks and planets, including HD 95086, HD 106906, and HD 133803. These objects are about 15-17 Myr old. The thermal emission from the debris disks constrains the locations of the dust. The dust is typically interior to the directly imaged planets in the systems. If additional planets reside in these systems, their locations are constrained by the positions of the dust belts. Many debris disk systems in Sco Cen appear to be two-belt systems. The gap between the belts in each system is a likely location for additional planets. The detection of planets in debris disk systems provide clues about the planet formation process, giving insights into where, when and how planets form.

Author(s): Hannah Jang-Condell⁸, Christine H Chen⁴, Tushar Mittal⁶, Erika Nesvold¹, Marc J Kuchner³, P Manoj⁵, Dan Watson⁷, Carey M Lisse²
Institution(s): 1. *Carnegie Institution for Science*, 2. *Johns Hopkins University Applied Physics Laboratory*, 3. *NASA Goddard Space Flight Center*, 4. *Space Telescope Science Institute*, 5. *Tata Institute of Fundamental Research*, 6. *University of California, Berkeley*, 7. *University of Rochester*, 8. *University of Wyoming*

105.04 – The survival of gas giant planets on wide orbits

It is not known whether gas giant planets on wide orbits form the same way as Jupiter or by fragmentation of gravitationally unstable discs. It has been suggested that giant planets that form on wide orbits in gravitationally unstable discs quickly migrate towards the central star. We simulate the migration of such planets including the effects of gas accretion onto the planet and radiative feedback from the planet, both of which have been ignored in previous studies. We show that a giant planet, which has formed in the outer

regions of a protostellar disc, initially migrates towards the central star while accreting gas from the disc. However, the planet eventually opens up a gap in the disc and the migration is essentially halted. At the same time, accretion-powered radiative feedback from the planet, significantly limits its mass growth, keeping it within the planetary mass regime (i.e. below the deuterium burning limit). Giant planets are therefore able to survive as planets (not higher-mass objects, i.e. brown dwarfs) on wide orbits, shaping the environment in which terrestrial planets that may harbour life form.

Author(s): Dimitris Stamatellos¹
Institution(s): 1. *University of Central Lancashire*

105.05 – Constraint of a planet mass from the depth and width of an observed gap on a protoplanetary disk

In a protoplanetary disk, a large planet is able to create the so-called disk gap, which is a low gas density region along the planet's orbit, due to the gravitational interaction between the disc and the planet. The gap formation induced by the giant planet is a possible mechanism to explain the formation of the so-called pre-transition disks with a ring gap structure. If the gap is created by the planet, the gap shape, i.e., the depth and width, would represent the mass and location of the planet. At the present stage, many pre-transition disks have been observed by e.g., ALMA and Subaru telescopes. It is important for us to examine what properties of the planet are constrained from the observed gap if the planet is in the gap.

We derived the relation between the depth of the observed gap and the planet mass in the gap based on the analytical model (Kanagawa et al. 2015a). This relation is a powerful tool to estimate the planet mass from the direct imaging of gaps in protoplanetary disks. We also applied this relation to the image of HL Tau' disk given by a part of the 2014 ALMA long baseline campaign and estimate the planet masses (Kanagawa et al 2015b). We also performed the numerical hydrodynamic simulation with the FARGO which is well-known code for the rotation disk, and found that the gap width becomes wider with a square root of the planet mass. Using this empirical relation for the gap width, we can also constrain the planet mass from the gap width. I'll talk about the relation between the gap depth, width and the planet, and the method for estimating the planet mass from the observed image of the disks.

Author(s): Kazuhiro Kanagawa², Takayuki Muto³, Hidekazu Tanaka², Takayuki Tanigawa⁴, Taku Takeuchi¹
Institution(s): 1. *Tokyo Institute of Technology*, 2. *ILTS, Hokkaido University*, 3. *Kogakuin university*, 4. *University of Occupational and Environmental Health*

105.06 – Using Protoplanetary Disks to Weigh the Youngest Stars and Constrain The Earliest Stages of Stellar Evolution

Mass is the fundamental property that determines the fate of a star. In particular, the masses of young stars are of great relevance to many astrophysical problems, including star and planet formation. We have developed a novel approach that combines spatially resolved sub-millimeter spectral line imaging and optical/near-infrared high resolution spectroscopy to derive the fundamental properties of a young star: mass, temperature, and radius. By applying our technique to a sample of pre-main sequence stars, we are mapping out a dynamically-calibrated Hertzsprung-Russell diagram for the express purpose of evaluating pre-main sequence evolutionary models. Looking forward, ALMA is poised to deliver precise stellar masses in statistically large quantities, enabling a meaningful survey of the fundamental properties of young stars.

Author(s): Ian Czekala¹, Sean Andrews¹, Eric Jensen², Keivan Stassun³, David W Latham¹, David Wilner¹, Guillermo Torres¹
Institution(s): 1. *Harvard-Smithsonian Center for Astrophysics*, 2. *Swarthmore College*, 3. *Vanderbilt University*

105.07 – Caught in the act: The quest for forming giant planets still embedded in their parent disk