

Swarthmore College

## Works

---

Senior Theses, Projects, and Awards

Student Scholarship

---

Spring 2021

### Ocean acidification modulates the impact of fluoxetine on larval behaviors of non-target organisms

Veronica A. Chua , '21

Follow this and additional works at: <https://works.swarthmore.edu/theses>



Part of the [Biology Commons](#)

---

#### Recommended Citation

Chua, Veronica A. , '21, "Ocean acidification modulates the impact of fluoxetine on larval behaviors of non-target organisms" (2021). *Senior Theses, Projects, and Awards*. 167.

<https://works.swarthmore.edu/theses/167>

This work is brought to you for free by Swarthmore College Libraries' Works. It has been accepted for inclusion in Senior Theses, Projects, and Awards by an authorized administrator of Works. For more information, please contact [myworks@swarthmore.edu](mailto:myworks@swarthmore.edu).

**Ocean acidification modulates the impact of fluoxetine on  
larval behaviors of non-target organisms**

by

Veronica Angeline Chua

Advisor: Kit Yu Karen Chan, Assistant Professor

Biology, Swarthmore College

A Thesis Submitted to the Biology Department  
in Partial Fulfillment of the Requirements of the Honors Program

May 2021, Swarthmore College

## Abstract

Emerging pollutants, such as pharmaceuticals from human waste, are continuously released into aquatic systems. Although pharmaceuticals alone can adversely impact marine organisms, the bioavailability of many pharmaceuticals are dependent on ambient physical conditions, like pH. As few studies have considered the interactive effects of pharmaceutical pollution and anthropogenic ocean acidification, this study investigated the behavioral response of larval sea urchins (*Heliocidaris crassispina*) and ascidians (*Styela plicata*) to environmentally-relevant concentrations of fluoxetine (10 and 100 ng L<sup>-1</sup>) under ambient (pH 8.0) and acidified conditions (pH 7.7). Larval ascidians reared at pH 8.0 exhibited swam in slower, more directed paths with increasing fluoxetine. Interestingly, this effect was absent at pH 7.7. On the other hand, I only observed independent effects of fluoxetine and acidification on urchin swimming behavior. My findings highlight the importance of using behavioral endpoints when assessing the realistic sub-lethal organismal and ecological impacts of anthropogenic stressors, and that considering differences in species traits may allow for the generation of more realistic predictions of the impact of emerging pollutants under future climate scenarios.

## 1. Introduction

<sup>1</sup>Increasing human interference of aquatic ecosystems continues to threaten marine biodiversity. Modern anthropogenic stressors range from ocean warming and acidification caused by increasing carbon emissions (Caldeira & Wickett 2005; Ricke & Caldeira 2014) to contamination with regulated pollutants (Förstner & Wittmann 1981; van Grinsven *et al.* 2012) and emerging pollutants with yet-to-be determined impacts (Deblonde *et al.* 2011; Geissen *et al.* 2015). Within the long list of emergent pollutants, many chemical compounds have the potential to interact with physical stressors to create non-linear impacts on marine organisms (Bryant *et al.* 1985; Stockdale *et al.* 2016; Almeida *et al.* 2018). Understanding the likelihood, magnitude, and mechanisms of such interactions are crucial for determining management goals and strategies (Boyd *et al.* 2018; Sarà *et al.* 2018).

---

<sup>1</sup> Data presented here is a part of larger collaboration with H.K. Lo from the Hong Kong University of Science and Technology. The resulting manuscript is in review in the journal *Science of the Total Environment*.

Environmental exposure to pharmaceuticals, a class of emergent pollutants, is of urgent concern due to their potential to adversely affect non-target organisms via ubiquitous biological pathways (Gaw *et al.* 2014; Fabbri & Franzellitti 2016). Most pharmaceuticals are only partially absorbed by target organisms (Monteiro & Boxall 2010), leaving trace concentrations in wastewater effluent that are not fully removed by treatment (Kot-Wasik *et al.* 2016). One such pharmaceutical is fluoxetine (FX), a selective serotonin reuptake inhibitor (SSRI) that increases signal transmission across serotonergic synapses and is the active compound in many antidepressant medications (Song *et al.* 1993). Fluoxetine HCl (tradename Prozac® or Sarafem®) is considered to be one of the most acutely toxic pharmaceuticals for non-target organisms, as evidenced by its low EC<sub>50</sub> and LC<sub>50</sub> values (Fent *et al.* 2006).

Given that serotonergic pathways are conserved across a wide array of taxonomic groups and are essential for regulating locomotion and development (Hen 1993), it is not surprising that FX exposure is associated with negative behavioral impacts. At concentrations up to 3.45 mg L<sup>-1</sup>, FX has been reported to decrease feeding in worms (Hird *et al.* 2016), decrease crawling speed in snails (Fong *et al.* 2015), and impair sand digging behavior in cuttlefish hatchlings (Di Poi *et al.* 2013). FX also increased locomotion in estuarine crabs (Peters *et al.* 2017) and increased photosensitivity and geotaxis in amphipods (Guler & Ford 2010). Apart from affecting behavior, serotonin also plays a crucial role in early embryogenesis (Buznikov *et al.* 2001). FX can also affect early juveniles and adults, as evidenced by delayed maturity in polychaetes (Méndez & Barata 2015) and induced spawning in mussels (Lazzara *et al.* 2012). At environmentally-relevant concentrations, zebrafish were able to biotransform FX into its

metabolite norfluoxetine, reducing the overall bioaccumulation of FX (Zindler *et al.* 2020). However, these studies did not account for the possible interaction between FX and other physical ocean conditions, such as temperature or pH.

FX is relatively stable in aquatic environments, with a half-life exceeding 100 days (Kwon & Armbrust 2006). Additionally, its high sorption coefficient suggests it can be readily transported and accumulated in soil and sediment (Yamamoto *et al.* 2005), increasing its ability to persist in aquatic ecosystems in direct contact with wastewater effluent, where FX concentrations can range from 12 to 540 ng L<sup>-1</sup> (Weston *et al.* 2001; Kolpin *et al.* 2002). For example, recent coastal measurements near the Asian megacity of Hong Kong revealed FX concentrations of up to 97.2 ng L<sup>-1</sup> (Lo 2020). Coastal regions are hotspots for pharmaceutical occurrence due to their proximity to human activities (Gulkowska *et al.* 2007; Singh *et al.* 2010; Fenet *et al.* 2014). Further, the rising usage of relatively stable pharmaceuticals like FX (Mojtabai & Olsson 2014), combined with the continuous discharge of wastewater into coastal ecosystems, can lead to the long-term “pseudo-persistence” of these contaminants, posing harmful effects to non-target organisms if left unregulated (Gelatti *et al.* 2013; Polverino *et al.* 2021).

The pH for maximizing FX bioavailability is alkaline at 13.7, where over 99.99% exists in its original form; however, below pH 5.6, more than 99.99% of FX is ionized (norfluoxetine), yielding the lowest bioavailability (Brooks *et al.* 2003). Pharmaceutical bioavailability is especially important to consider in the context of ocean acidification (OA), where increasing anthropogenic carbon emissions drive changes in ocean carbonate chemistry, resulting in a predicted pH drop of 0.2 to 0.4 units by the end of the century (Caldeira & Wickett 2005). In addition, coastal regions experience frequent and

large fluctuations in pH that can reach end-of-century pH even at present-day (Hofmann *et al.* 2011; Pecquet *et al.* 2017). For instance, the Pearl River Delta, which includes Hong Kong, experienced a pH drop from 8.2 to 7.7 in the last two decades (Wang *et al.* 2016). Since the bioavailability—and thus potential impact on non-target organisms—of pharmaceuticals, like FX, is modulated by ambient pH, studying the interactive effects of these two anthropogenic stressors will help inform future performance predictions and the management of coastal communities.

To investigate the hypothesis that FX-induced biological impacts would be mitigated by OA due to decreased bioavailability, I exposed the sea urchin *Heliocidaris crassispina* and the ascidian *Styela plicata* to environmentally-relevant concentrations of FX under near-future OA conditions. Sea urchins and ascidians are two ecologically and commercially important model organisms generally found in coastal regions. *H. crassispina* are grazers commonly found in the shallow, rocky habitats along the coasts of East Asia (Chiu 1990; Pearse 2006; Agatsuma & Lawrence 2013). Sea urchins serve as prime model organisms for developmental or neurobiological studies (Epel *et al.* 2006; Aluigi *et al.* 2010, 2012). For example, the purple sea urchin *Strongylocentrotus purpuratus*, whose genome has been fully sequenced (Burke *et al.* 2006), has homologs for serotonin transporters (SERT), as well as Type 1, 2, and 7 serotonin receptors and the  $\gamma$ -aminobutyric acid type A (GABA<sub>A</sub>) receptor, all of which are involved in FX target pathways (Tunnickliff *et al.* 1999; Fabbri & Franzellitti 2016).

*S. plicata* is a fouling species with a wide-ranging distribution (Pineda *et al.* 2011; Aldred & Clare 2014), often found in coastal waters with large environmental fluctuations and human disturbance (Naranjo *et al.* 1996). Ascidians in general are

evolutionarily close to vertebrates with a relatively simple central nervous system (Lemaire *et al.* 2008). The ascidian *Ciona intestinalis* also has a fully sequenced genome (Dehal *et al.* 2002) and a well-documented developmental cycle (Satoh 1994), making ascidians prime model organisms for studies in neurobiology (Dahlberg *et al.* 2009), developmental and evolutionary biology (Corbo *et al.* 2001), and ecotoxicology (Gallo & Tosti 2015). Previous studies have observed SERT-expressing coronet cells in the *Ciona* sensory vesicle, as well as genes related to the GABA<sub>A</sub> receptor and putative serotonin receptors (Okamura *et al.* 2005; Horie *et al.* 2009; Razy-Krajka *et al.* 2012).

Planktonic larvae are generally more susceptible to environmental stressors than other life stages (Pechenik 1999). Additionally, comparing the larval stages of two non-target organisms provides an informative framework to investigate species-specific sensitivity to combined climate change stressors (Byrne & Przeslawski 2013). Unlike ascidians, many sea urchin larvae are highly calcifying. Calcifiers are suggested to be highly sensitive to OA due to the dissolution of calcified structures (Bednaršek *et al.* 2012), increased metabolic costs of calcification (Stumpp *et al.* 2012), and/or decreased availability of carbonate at low pH (Hofmann *et al.* 2010). Urchin larvae beat cilia to regulate their position in the water column (Grünbaum & Strathmann 2003), whereas ascidian tadpoles are undulatory muscular swimmers (McHenry 2005). Urchin larvae also use these ciliary bands to feed (Strathmann & Grünbaum 2006) while *S. plicata* larvae are non-feeding (David *et al.* 2010), possibly suggesting differential exposure opportunities between the two species.

In the present study, I analyzed the swimming behavior of *H. crassispina* and *S. plicata* larvae to test if combined exposure to OA and FX interactively affect organismal

performance. While most ecotoxicological assays rely on mortality and other physiological markers, behavioral endpoints are integrative, whole-organism responses that provide nuanced information about the sublethal impacts of environmentally-relevant stressors (Gerhardt 2007; Ford *et al.* 2021). Additionally, alterations in larval swimming behavior hold strong implications for dispersal dynamics and population connectivity (Cowen & Sponaugle 2009). My findings reveal multiple behavioral impacts of the non-linear interaction between OA and FX exposure in ascidian larval swimming, as well as altered urchin behavior resulting from independent effects of the two stressors.

## **2. Materials and methods**

### **2.1 Animal collection and spawning**

Adult sea urchins (*Heliocidaris crassispina*) were procured from a fish farm in Hong Kong (22.39° N, 114.11° E) and transported to the HKUST Coastal Marine Laboratory. Adults were kept in  $21 \pm 1^\circ\text{C}$ , 200  $\mu\text{m}$ -filtered seawater upon arrival and fed a diet of dried kelp. After three months, spawning was induced by injecting 0.5-1 mL of 0.5 M KCl into the intracoelomic cavity (Strathmann 1987). Eggs from three females were collected in separate beakers with MBL artificial seawater (ASW, Cavanaugh 1956,  $\text{pH}_T = 8.007$ ,  $20.1^\circ\text{C}$ ), followed by washing through a 125- $\mu\text{m}$  mesh to remove large debris. Sperm from three males were collected dry on ice before use. Eggs were fertilized with a sperm concentration of  $1000 \text{ mL}^{-1}$  (3 females x 3 males). Fertilization success (> 95%) was confirmed under a microscope, with the fertilization envelope observable at 5 minutes post-fertilization (mpf). Fertilized eggs were washed with ASW to remove excess sperm, then mixed and allocated within 30 mpf to 1.5 L glass jars (2.5 individuals



mL<sup>-1</sup>) at their respective treatments ( $20 \pm 1$  °C).

Adult *Styela plicata* were collected by detaching them from ropes on a mariculture raft (Yung Shue Au, Hong Kong; 22.39° N, 114.11° E) and also transported to the HKUST Coastal Marine Laboratory. Prior to dissection, *S. plicata* were maintained at  $21 \pm 1$  °C for one week using free-flowing seawater. Gametes were collected by cutting through the tunics of one individual to expose the hermaphroditic gonads. Oocytes were stripped from the gonad and immediately rinsed with ASW (pH<sub>T</sub> = 8.10, 19.6°C) through a 250-µm, then 45-µm mesh filter to remove large particles. Sperm were collected from another 3 individuals by stripping the whitish spermiduct and testes, after which they were observed under a microscope to confirm active swimming. Gametes were combined at a 4:1 sperm-to-oocyte ratio. At 5 mpf, eggs were washed with ASW three times before filtering through a 45-µm mesh, then allocated into 400-mL jars (2.5 individuals mL<sup>-1</sup>) at their corresponding treatments within 15 mpf ( $20 \pm 1$  °C).

## 2.2 Experimental design

Larvae were exposed to a full factorial design of three nominal FX concentrations (control = 0 ng L<sup>-1</sup>; low = 10 ng L<sup>-1</sup>; high = 100 ng L<sup>-1</sup>) and two pH levels (ambient control = 8.0; OA = 7.7), yielding 6 treatments with 3 replicates per treatment. The low and high FX concentrations reflect present-day pre- and post-typhoon measurements in Hong Kong (Lo, 2020). Notably, the high FX value is conservative relative to the concerning environmental prediction of 370 ng L<sup>-1</sup> in UK drinking water (Webb 2004). The pH of the OA condition reflects the predicted reduction in global surface ocean pH

by the end of the century (Caldeira & Wickett 2005), and is observed within present-day pH fluctuations in Hong Kong waters (Wang *et al.* 2016; Pecquet *et al.* 2017).

ASW was used throughout the entire experiment to ensure that the nominal FX concentrations would not be affected by potential traces of FX in natural seawater. Prepared ASW was vigorously aerated overnight with compressed air filtered through a 0.22  $\mu\text{m}$  membrane filter before use. Urchin larvae were fed with 5000 cells  $\text{mL}^{-1}$  *Rhodomonas salina*. Algal concentration was verified with a Coulter Counter (Multisizer 4e, Beckman Coulter, United States), and a complete water change was performed every 2 days. Due to the short-lived and non-feeding larval duration of *S. plicata*, ascidian larvae remained undisturbed until behavioral analysis.

### **2.3 Seawater carbonate and fluoxetine chemistry**

Filtered ASW in each jar was continuously and gently bubbled with air passed through a 0.22- $\mu\text{m}$  filter. Jars at nominal  $\text{pH}_T$  7.7 were connected to pH computers (Aqua Medic, Germany) to control pH through the addition of pure  $\text{CO}_2$ . Temperature, voltage, and  $\text{pH}_{\text{NBS}}$  were monitored daily with a Metrohm 913 pH meter and unitrode with Pt 1000 (Herisau, Switzerland). Salinity was measured with a handheld refractometer. To convert measured pH to total scale pH ( $\text{pH}_T$ ), daily measurements were made on a Tris/HCl buffer solution (salinity = 33) provided by the Dickson Lab.

Total alkalinity ( $A_T$ ) and dissolved inorganic carbon (DIC) were measured for each jar and the newly prepared ASW of corresponding pH during each water change. Water samples for  $A_T$  measurement were filtered through a 0.22  $\mu\text{m}$  syringe filter, while DIC samples were collected in air-free vials without filtration. All water samples were

stored at 4 °C before analysis.  $A_T$  was quantified using a computer-driven titrator (905 Titrand, Metrohm, Switzerland) mounted with a glass electrode (Unitrode with Pt1000, Metrohm, Switzerland). DIC was quantified by LiCor Mass Spectrometry (AS-C3, Apollo Technology Solutions LLC, USA). Calibration was performed using the standard seawater provided by the Dickson Lab (Batch 151). Carbonate system parameters ( $pCO_2$ ,  $\Omega_{ar}$ , and  $\Omega_{ca}$ ) were calculated from these measurements in CO<sub>2</sub>SYS using the dissociation constants from Mehrbach *et al.* (1973) as refitted by Dickson & Millero (1987).

Standard fluoxetine HCl (PHR1394, Sigma-Aldrich) was used to prepare a working standard of 40  $\mu\text{g L}^{-1}$  and stored in darkness at 4 °C. The nominal concentration of working standard was verified with liquid chromatography with electron spray and tandem mass spectrometry (LC-ES-MS/MS) using a modified version of the methods described by Chu & Metcalfe (2007). Given the quantification of FX at the  $\text{ng L}^{-1}$  level requires at least one liter of water sample, an end-point FX concentration reading from the rearing jar (1.5 L volume) could not be obtained. However, since identical glassware were used and the same amount of algal cells (>300  $\mu\text{g}$  dry mass) were added across all treatments, the loss, if any, would be comparable across all treatments.

## **2.4 Swimming behavior in response to OA and FX exposure**

Larval swimming behavior was recorded at 7 dpf for urchins and 16 hpf for ascidians. Duplicate 10 mL subsamples ( $\approx 15\text{-}20$  larvae) from each replicate jar were each placed in a well of a 6-well plate. After two hours of acclimation, the plate was moved onto the base of a dissecting microscope (MZ6, Leica) mounted with an eyepiece camera (AM4023X, Dino-Lite, Taiwan). The field of view covered the only center portion of the

well to avoid wall effect. After an additional 15 minutes, each well was videotaped for 3 min at 15 fps (at least 7 individuals in view). The process was repeated for all 6 treatments. Formalin-fixed larvae were also filmed for 3 min to account for potential drifting. Room temperature was maintained at  $20 \pm 1$  °C, and water temperature during recording was monitored with a thermocouple.

Videos were processed in FOSICA (Wallingford Imaging, WA), first by removing background and thresholding for size and brightness, then converting the positions of moving particles into pixel coordinates. I used the in-house MATLAB program Tracker3D (Chan *et al.* 2011) for path assembly (Fig. 1a, b). Larval swimming speed was measured by gross and net speed. Gross speed was calculated by applying a smoothing spline to the paths to remove frame rate noise and taking the time derivatives of the splines. Net speed was calculated as the displacement between the first and last coordinates over time. I also computed net-to-gross displacement ratio (NGDR) and mean squared displacement (MSD). NGDR was calculated by dividing the net displacement of a path by the gross displacement. This metric can range from 0 to 1, with low values indicating convoluted trajectories and high values indicating straightforward trajectories (True *et al.* 2015). Similarly, MSD was calculated as:

$$MSD_{path} = \frac{1}{N-j} \sum_{i=1+j}^N |r_i - r_{i-j}|^2$$

where  $N$  is the number of datapoints collected from the path,  $j$  is the chosen time interval, and  $r$  is the physical position of the larva. The MSD of each path was interpreted as the distance explored through Brownian motion, indicating the level of the trajectory's convolution. All swimming metrics were calculated in MATLAB.

Beyond speed and trajectory, I also investigated the effect of combined FX and

OA on discrete larval behaviors. Orientations of urchin larvae (ventral, dorsal, animal, vegetal, left, or right; Fig. 1c) were recorded for all visible larvae during the first, middle, and last frame of each video. Orientations were observed as the side facing the camera (bird's eye view from above the water surface) in a given frame. Ethograms of ascidian movement were created by recording which behavior was being executed for the first 900 frames (1 min) of all videos. Ascidian movement was placed into one of six categories (Fig. 1d). "Still" ascidian larvae did not move their tails or trunks between the current frame and the next frame. "Swimming" larvae moved their tails in the classic undulatory motion, resulting in trunk-directed displacement between frames. "Cruising" larvae also exhibited trunk-directed displacement, but moved without any visible tail manipulation. Larvae were considered "spasming" if there was observable tail movement compared to the previous and/or next frame, but no trunk-directed displacement. "Twitching" larvae exhibited unilateral contraction and relaxation of the tail (i.e., a straight larva bending into a C-shape, then returning to its original position). Finally, "spinning" larvae rapidly contracted their tails resulting in clockwise or counterclockwise rotation around the base of the trunk. As the field of view only excluded the edges of the well, individuals were visible in the field of view for varying amounts of time. Thus, behaviors were calculated as the percent of total time spent by all individuals in a well conducting each behavior.

## **2.5 Statistical analysis**

To determine the effect of OA and FX exposure on larval swimming behaviors (speeds, trajectory, frequency of urchin orientation, and percent time ascidian larvae spent conducting categorical behaviors), I employed a 2-way ANOVA with pH and FX

as fixed factors. All data were checked with a Shapiro-Wilks test for normality and Levene's test for homogeneity of variance. Ascidian ethogram data were arcsine square root transformed before statistical analysis. In the cases where data did not fit normality and homogeneity assumptions, I analyzed the data using an aligned rank transform, non-parametric ANOVA (i.e., ARTool; Wobbrock *et al.* 2011) and significant *post hoc* effects were analyzed with a difference-of-differences contrast. The level of significance was set to  $p < 0.05$ . All statistical tests were performed in R (R Core Team Version 4.0.2).

### **3. Results**

#### **3.1 Seawater carbonate and fluoxetine chemistry**

Rearing conditions for urchin larvae at ambient pH 8.0 and in OA pH 7.7 were stable over the course of the experiment for both urchins and ascidians ( $F_{1,58} = 0.66$ ,  $p > 0.5$  and  $F_{1,62} < 10$ ,  $p > 0.05$ , respectively; Table 1). pH in the ambient control was significantly different from pH in the OA condition ( $F_{1,58} = 972$ ,  $p < 0.0001$  and  $F_{1,62} < 200$ ,  $p < 0.05$  for urchins and ascidians, respectively). The average measured FX concentration in the working standard was  $39.15 \mu\text{g L}^{-1}$  ( $\approx 97.9\%$  of  $40 \mu\text{g L}^{-1}$ ) for the urchin experiment and  $40.09 \mu\text{g L}^{-1}$  ( $\approx 100.23\%$  of  $40 \mu\text{g L}^{-1}$ ) for the ascidian experiment.

#### **3.2 Larval urchin swimming behavior**

OA alone ( $F_{1,19} = 11.8$ ,  $p < 0.01$ ; Fig. 2a) significantly slowed urchin gross swimming speed at all FX concentrations. FX exposure alone marginally affected gross speed ( $F_{2,19} = 3.85$ ,  $p = 0.051$ ), with the low concentration reducing gross speed more than control and high concentrations. Neither net speed nor NGDR of larval urchins was

significantly affected by OA, FX, or their interaction (Fig. 2b, c). However, I did observe a slight increase in NGDR when exposed to FX and OA independently. MSD of larvae in pH 8.0 was significantly higher than larvae reared in OA ( $F_{1,19} = 7.44, p < 0.05$ ) regardless of FX exposure (Fig. 2d). Additionally, I observed a significant interaction of OA and FX concentration on the frequency of left-side-up larvae ( $F_{2,20} = 8.71, p < 0.05$ ; Fig. 3a). At low FX concentrations, there were less left-side-up larvae in reduced pH; this pattern was reversed at high FX concentrations ( $F_{1,12} = 8.71, p < 0.05$ ). Larvae were also oriented dorsal-side-up more often in increased FX concentrations, though this effect was only marginally significant ( $F_{2,20} = 3.72, p = 0.056$ ; Fig. 3b). There were no significant effects of pH, FX, or their interaction on any of the other four orientations measured.

### **3.2 Larval ascidian swimming behavior**

Both OA ( $F_{1,11} = 13.0, p < 0.005$ ) and FX ( $F_{2,11} = 5.03, p < 0.05$ ) independently affected median ascidian net speed. I also observed a significant interactive effect ( $F_{2,11} = 12.3, p < 0.005$ ), wherein larval ascidians reared in ambient pH had steeper decreases in net speed at low ( $F_{1,11} = 9.97, p < 0.05$ ) and high FX ( $F_{1,11} = 23.7, p < 0.005$ ) from control FX (Fig. 4b). However, ascidian larvae exposed to OA exhibited no significant differences in net speed across FX. In other words, ascidian net speeds decreased with increasing FX at ambient pH, whereas this trend was absent in OA. While not statistically significant, this pattern is reflected in the NGDR (Fig. 4d). Additionally, there was no significant effect of pH, FX, or their interaction on gross speeds or MSD (Fig 4a, c).

I did not observe a significant difference in the percent of overall time spent by ascidians conducting any of the 6 categorical movement behaviors. However, analysis of

behavior combinations revealed that ambient pH ascidians reared in control FX ( $14 \pm 7.1\%$ ; mean  $\pm$  SD) spent a longer, but non-significant, proportion of time conducting translating behaviors (“swim” + “cruise”) than those at high FX ( $2.1 \pm 1.3\%$ ; mean  $\pm$  SD) (Fig. 3c). Concordantly, ascidians at ambient pH spent more time being still at high FX ( $96 \pm 0.4\%$ ; mean  $\pm$  SD) than at control FX ( $84 \pm 6.1\%$ ; mean  $\pm$  SD). However, this FX-dependent pattern is absent under OA. There is a significant interaction between pH and FX on the number of individuals that remained still for the entire recording ( $F_{2,12} = 4.94$ ,  $p < 0.05$ ), the number of long-term still individuals peaks in low FX ( $F_{1,12} = 6.59$ ,  $p < 0.05$ ) under OA but in high FX ( $F_{1,12} = 8.14$ ,  $p < 0.05$ ) in control (Fig. 3d).

#### 4. Discussion

While antagonistic, synergistic, and additive interactions have been reported between pH and other pollutants (Freitas *et al.* 2016; Dorey *et al.* 2018), little is known about the extent and patterns of interactive effects between pharmaceutical pollution and OA on non-target marine organisms. I investigated whether FX-induced impacts are modulated by pH levels relevant to end-of-century OA predictions. By exposing early developmental stages of the urchin *H. crassispina* and the ascidian *S. plicata* to environmentally-relevant pH and FX levels, this study reveals species-specific non-linear impacts on behavior. Rearing ascidians in OA mitigated the reductions in net speed induced by FX exposure; however, OA and FX acted independently on the speed and trajectory of urchin larvae. Thus, recreating realistic field conditions is crucial for predicting the non-linear and species-specific impacts of emergent pollutants on non-target marine organisms in future ocean scenarios.



#### 4.1 OA modulates FX-induced impacts on ascidian swimming

I observed an interactive effect between pH and FX exposure on larval ascidian swimming behaviors. At ambient pH 8.0, net speeds, NGDR, and the time spent exhibiting directed movement decreased with increasing FX concentration; however, this effect was not observed in OA conditions. This pH-dependent modulation of FX impacts could be a result of reduced bioavailability of FX under reduced pH. Within the narrow range of pH investigated, pKa predictions show that at pH 8.3, only 3% of FX remains unionized, further reducing to 0.78% at pH 7.7 (Lo 2020). Previous studies suggest that norfluoxetine is more soluble and has a lower chance of bioaccumulation and reduced bioavailability (Brooks *et al.* 2003). Indeed, reduced FX bioavailability has been observed at pH 7.0 compared to pH 8.0 and 9.0, resulting in reduced acute toxicity and bioaccumulation in Japanese medaka tissues (Nakamura *et al.* 2008).

Another possible driver for pH-dependent responses to FX exposure could be a shift in energy allocation due to increased physiological stress. A companion study to this thesis found that *S. plicata* larvae reared in ambient pH 8.0 had higher levels of DNA damage with increasing FX exposure (Lo 2020). Ascidians rely on apoptosis to defend against pollutant-induced oxidative stress (Barbosa *et al.* 2018). Given that apoptosis is energy-dependent (Eguchi *et al.* 1997; Elmore 2007), it may be the case that at higher levels of stress (i.e., high concentrations of bioavailable FX), ascidians shift energy allocation to prioritize maintenance over swimming, which can also be energetically expensive (Bennett & Marshall 2005). Such trade-offs have been observed in other aquatic invertebrates under environmental stress (Sokolova *et al.* 2012). Relatedly, *S. plicata* larvae did not experience a change in respiration rate when exposed to OA and

FX conditions (Lo 2020). This observation is not surprising given that ascidian larvae are non-feeding and must rely on limited maternal provisions to complete their larval duration (Berrill & Watson 1935).

The observed reduction in net speed and NGDR suggest that in ambient pH, ascidians swim with more convoluted paths than in high FX concentrations. Razy-Krajka *et al.* (2012) showed that high concentrations of FX decreased spontaneous swimming of *Ciona* by acting on the SERT-containing cells in the sensory vesicle, which are involved in sensoreception. However, this study only measured tail beat frequency, and did not account for other metrics of swimming kinematics (e.g. differences in yaw), which may reveal effects of FX on swimming trajectory (McHenry 2005). Additionally, the exact mechanism behind FX action is still unclear, as SERT-containing cells in ascidians are multi-functional and may facilitate transport of other neurotransmitters like dopamine (Razy-Krajka *et al.* 2012). Rudolf *et al.* (2019) also showed that modafinil, a selective dopamine transporter inhibitor, increased swimming speed of *Ciona*. Similarly, dexmedetomidine, an ADR $\alpha$ 2 dopamine receptor agonist, also increased spontaneous swimming in ascidian larvae (Razy-Krajka *et al.* 2012).

Interestingly, ascidians exposed to OA exhibited relatively low net speeds and NGDR regardless of FX concentration. OA has been hypothesized to reverse GABAergic signaling by affecting ion gradients (Nilsson *et al.* 2012). Inhibition of GABAergic signaling has been observed to increase the frequency and duration of spontaneous swimming events in *Ciona* larvae (Brown *et al.* 2005); however, it has also been observed to reduce negative geotaxis in response to light (Bostwick *et al.* 2020). The latter behavior refers to the “light-off response”, where the dimming of visible light

induces negative geotaxis and a rapid increase in swimming, followed by recovery after visible light exposure resumes (Nakagawa *et al.* 1999; Tsuda *et al.* 2003; Zega *et al.* 2006). While I demonstrated that OA and FX affect ascidian swimming behavior in an interactive manner, further studies on the complex sensorimotor mechanisms involved in the light-off response will provide insight into how combined climate change stressors affect neuroethology in a more ecologically relevant context.

One possible approach is to analyze ascidian swimming in response to light change, using cuvettes that allow for geotactic movement (Fig. 5). As my results provide insight on the impact of rearing ascidian embryos in combined stressors, acute exposure assays may help tease apart the effect of developmental abnormalities and resulting behavior. Preliminary data suggest that ascidian larvae exposed to combined OA and FX still exhibit increased swimming responses to light dimming; however, the presence of an interaction requires further in-depth analysis (Chua, unpub. data). I hypothesize that OA may antagonistically interact with FX due to increased serotonergic signaling from the sensory vesicle, depending on where GABA reversal acts on the ascidian neural system. Results supporting this hypothesis would suggest that the change in behaviors associated with OA- and FX-exposure has a neurological basis.

#### **4.2 OA and FX independently affect larval urchin swimming**

In contrast with the interactive effect on ascidian behavior, I observed an FX-independent reduction in gross swimming speed and MSD of urchin larvae in OA compared to those at ambient pH. Other echinoid larvae maintained swimming speed under acidified conditions (e.g. *Dendraster excentricus*, Chan *et al.* 2011;

*Strongylocentrotus droebachiensis*, Chan *et al.* 2015a; *S. purpurtaus*, Chan *et al.* 2015b). However, brittle star larvae swim slower at OA-relevant pH levels (Chan *et al.* 2015b), suggesting interspecific variation in swimming responses to acidification within echinoderms. The variation across studies may also be explained by differences in observation approach. Earlier studies focus on movement along the X-Z plane (vertical swimming), while the current study observed swimming in shallow (80 mm deep) well plates through the X-Y plane (horizontal swimming). For example, larval *D. excentricus* exposed to reductions in temperature experience reductions in horizontal speed to compensate for the maintenance of vertical movement (Chan & Grünbaum 2010).

Changes in larval morphology induced by pH stress could have also contributed to the observed difference in speeds. A companion study to this thesis found that larval bodies of *H. crassispina* were longer overall at ambient pH than in OA (Lo *et al.* in review). Smaller larvae may have relatively shorter ciliated bands to generate lift, thus reducing speed (Grünbaum & Strathmann 2003). Acidified echinoderm larvae were observed to have shorter arms and more U-shaped bodies compared to control, which affected swimming biomechanics, such as through increased stability in vertical shear (Chan *et al.* 2011; Chan & Tong 2020). We also observed changes in larval urchin arm symmetry under OA (Lo *et al.* in review), which may affect the larval weight distribution, and hence, swimming (Chan *et al.* 2011). Variation in arm symmetry may also explain the observed differences in left-side up orientation across pH treatments.

Aside from morphological differences, behavioral change could also be induced by low pH, such as through the hypothesized reversal of GABAergic signals in OA may modulate ciliary beating in urchin larvae (Kato *et al.* 2013). Previous studies have also

reported that larval *D. excentricus* avoided unfavorable environments, e.g., low pH (Maboloc *et al.* 2020) or high temperature (Chan & Grünbaum 2010). Here, larvae exposed to FX have a higher incidence of dorsal-side-up orientation at pH 8.0 than at 7.7. Though this difference is not statistically significant, increased downward swimming could be interpreted as an escape response. Given that I analyzed larval swimming in shallow dishes, future observations using free-swimming larvae in a larger volume may provide more insight on the implications of my observed changes in the field (Roy *et al.* 2012; Fuchs & Gerbi 2016).

I also observed that urchin larval gross speed was the slowest at low FX concentrations compared to control and high concentrations. Interestingly, I only observed a marginally significant effect of FX on larval gross speed, despite the regulatory role of serotonin in ciliary beating of sea urchin larvae (Mogami *et al.* 1992; Wada *et al.* 1997). While the negative effects of FX exposure on locomotion are well-documented in other organisms, these studies were conducted at concentrations several orders of magnitude higher than the current study (Hamilton *et al.* 2016; Ford *et al.* 2018). Further, it is possible for FX-induced behavioral changes to manifest over longer time scales than used in the current study, especially at lower concentrations (Di Poi *et al.* 2014; Peters & Granek 2016). Conducting whole life cycle analyses with environmentally-relevant FX levels, combined with future ocean conditions, would allow for more accurate ecological predictions of organismal responses.

#### **4.3 Interspecific variations in OA and FX responses**

Overall, FX impacts on larval behavior can be modulated by reduced pH, though

this effect is clearer in ascidians compared to urchins. This variation in impact may be due to species-specific traits. For example, the ascidian larval duration is limited by maternal provisions, and thus may be more pressed to reallocate energy and prioritize essential processes than urchin larvae, which may be able to meet energetic demands through food intake (Sokolova *et al.* 2012). Differential sensitivities to FX exposure have been observed in the polychaete *Hediste diversicolor* depending on feeding mode (Hird *et al.* 2016). Rather than the predicted modulation effect, I observed independent effects of OA and FX on larval urchin swimming. FX inhibits urchin toxicant defenses, e.g., expression of efflux transporters and cytochrome P450 (Lynch & Price 2007; Smith *et al.* 2012). However, reduced transcription of toxicant defense genes in OA (Todgham & Hofmann 2009) may be paired with the hypothesized reduced bioavailability of FX in OA, resulting in a similar extent of FX impact across different pH levels.

Additionally, species-specific differences in the neurophysiological control of swimming may help explain the observed variation in responses. The read-across hypothesis suggests that pharmaceuticals will affect non-target organisms as long as the molecular target is conserved (Rand-Weaver *et al.* 2013), though they may have different outcomes depending on the pathway the molecular target is involved in (Hird *et al.* 2016). SERT and other serotonergic receptors are involved in the swimming pathways of both urchins and ascidians; however, while serotonergic signaling plays a role in ciliary manipulation in urchin larvae, SERT is mainly involved in sensory reception in ascidian larvae. Further, while I did not observe a significant interaction of FX and OA on urchin swimming speed, a companion study found interactive effects on other physiological and

morphological metrics in urchin larvae (Lo *et al.* in review), supporting species-specific modes of action of FX.

## **5. Conclusion**

Exposure to environmentally-relevant levels of FX and pH altered behavioral responses in early life stages of the urchin *H. crassispina* and the ascidian *S. plicata*. However, while the two stressors independently impacted larval urchin responses, exposure to OA antagonistically modulated FX-induced impacts on larval ascidians. Changes in larval swimming behavior have implications for dispersal dynamics (Minchinton & Scheibling 1991) and survivorship (Chan & Grünbaum 2010; Chan *et al.* 2018). Previous research suggests that coastal organisms like ascidians and urchins may be more resilient to OA due to pre-exposure to low pH waters (Hofmann *et al.* 2010). My observations thus highlight the importance of using behavioral metrics as a sensitive tool when investigating sub-lethal impacts of climate change stressors. Future research should emphasize the use of multiple stressors, such as pharmaceutical mixtures (Backhaus 2014), as well as environmentally-relevant stressor levels to more accurately predict organismal responses to future ocean environments.

## **Acknowledgements**

This work was supported by the Research Grant Council of Hong Kong [Project No: 26102515 and 16127816] to Dr. Karen Chan. Data collection was conducted by Dr. H.K. Abby Lo; I thank D. Tong, H. Tso, C. Knox, E. Wong, Dr. Y.K. Tam for their assistance in the HKUST lab. I would also like to thank Darren Chen, Emily Branam, Pempho Moyo, Isaac Satz, Ismail Can, and the rest of my friends for their emotional support. I also would not have been able to complete this work without the love and support of my parents and my sisters, Alex and Jabee. Finally, I would like to acknowledge Dr. Karen Chan for her brilliant mentorship and guidance throughout the duration of this project.




## Tables


**Table 1.** Carbonate chemistry parameters of artificial seawater used to rear larval urchins (7 dpf) and ascidians (16 hpf). Temperature, total scale pH ( $pH_T$ ), and total alkalinity ( $A_T$ ) were measured, and dissolved inorganic carbon (DIC), partial pressure of CO<sub>2</sub> ( $pCO_2$ ), and aragonite and calcite saturation were calculated in CO<sub>2</sub>SYS. Values are reported as mean  $\pm$  SD.

	Measured parameters				Calculated parameters			
	Nominal pH	Temp (°C)	$pH_T$	$A_T$ ( $\mu\text{mol kg}^{-1}$ )	DIC ( $\mu\text{mol kg}^{-1}$ )	$pCO_2$ ( $\mu\text{atm}$ )	$\Omega_{ca}$	$\Omega_{ar}$
Urchin	8.0	19.5 $\pm$ 0.1	8.09 $\pm$ 0.03	2143 $\pm$ 8.2	2092 $\pm$ 37.6	375 $\pm$ 111	1.9 $\pm$ 0.6	1.3 $\pm$ 0.4
expt.	7.7	19.6 $\pm$ 0.1	7.70 $\pm$ 0.07	2166 $\pm$ 7.3	2169 $\pm$ 20.9	1181 $\pm$ 417	0.7 $\pm$ 0.2	0.5 $\pm$ 0.2
Ascidian	8.0	19.7 $\pm$ 0.2	8.11 $\pm$ 0.12	2181 $\pm$ 52.2	N.A.	349.6 $\pm$ 46	4.6 $\pm$ 1.0	2.9 $\pm$ 0.6
expt.	7.7	19.6 $\pm$ 0.2	7.76 $\pm$ .04	2178 $\pm$ 17.8	N.A.	822.2 $\pm$ 40	2.3 $\pm$ 0.2	1.4 $\pm$ 0.1


## Figures




**Figure 1.** Larval swimming behavior of *H. crassispina* urchins (7 dpf) and *S. plicata* ascidians (16 hpf) was quantified by calculating gross and net speeds, trajectory metrics net-to-gross displacement ratio (NGDR) and mean squared displacement (MSD), as well as discrete movement behaviors. **(A)** Diagram depicting a relatively convoluted path, characterized by low NGDR and high MSD. **(B)** Diagram depicting a relatively straightforward path, characterized by high NGDR and low MSD. Solid black lines indicate total distance traveled as used to calculate gross speed. Dashed lines indicate net displacement traveled as used to calculate net speed. **(C)** A 3D model of a pluteus larva at six discrete urchin larvae swimming orientations, which were used as a proxy to determine swimming direction. **(D)** Ascidian swimming behavior was broken down into six discrete movement categories. Red stop symbols represent no change in displacement over time. Heavy blue arrows represent change in displacement over time (i.e. “translating movement”).




**Figure 2.** Ocean acidification affects larval urchin swimming behavior. **(A)** Gross speed, **(B)** net speed, **(C)** net-to-gross displacement ratio (NGDR), and **(D)** mean squared displacement (MSD) were used to quantify the effect of ocean acidification (pH 7.7; red boxplots) and increasing fluoxetine concentrations compared to control pH 8.0 (blue boxplots). **(E)** Representative urchin paths at pH 8.0, characterized by high MSD and high gross speed. **(F)** Representative urchin paths at pH 7.7, characterized by low MSD and low gross speed. Letters indicate significantly different groups ( $p < 0.05$ ).



**Figure 3.** Ocean acidification and fluoxetine interact to affect larval urchin and ascidian behavior. **(A)** The proportion of larval urchins observed at left and **(B)** dorsal side-up orientations were recorded in response to acidification (pH 7.7; red boxplots) and increasing fluoxetine exposure compared to control pH 8.0 (blue boxplots). **(C)** The proportion of time (frames) larval ascidians spent conducting translating behaviors (swim + cruise; see Fig. 1) after exposure to acidification and increasing fluoxetine. **(D)** The number of ascidian larvae counted as completely still throughout the entire duration of the video (~3 min), weighted by the number of frames. Letters represent significantly different differences between groups ( $p < 0.05$ ), with subsequent numbers representing the groups being compared.



**Figure 4.** Ocean acidification mitigates fluoxetine-induced impacts on larval ascidian swimming behavior. **(A)** Gross speed, **(B)** net speed, **(C)** net-to-gross displacement ratio (NGDR), and **(D)** mean squared displacement (MSD) were used to quantify the effect of ocean acidification (pH 7.7; red boxplots) and increasing fluoxetine concentrations compared to control pH 8.0 (blue boxplots). **(E)** Representative urchin paths at pH 8.0 without fluoxetine, characterized by relatively high net speed and NGDR. **(F)** Representative urchin paths at pH 7.7 and 8.0 with fluoxetine, characterized by relatively low net speed and NGDR. Letters indicate significantly different differences between groups ( $p < 0.05$ ).



**Figure 5.** Video analysis of larval ascidian geotactic response to light dimming (i.e., “light-off” or “shadow” response). Two cameras fitted with 35 mm Nikon lenses recorded larval ascidians acutely exposed to combined fluoxetine (0, 10, and 100 ng L<sup>-1</sup>) and pH conditions (pH 8.0 and 7.7). Vertical cuvettes (white arrows) were used to allow for geotactic response to dimming of soft LED light (red arrow; ≈10 to 0 ppf). Cuvettes were constantly illuminated by IR light (yellow arrow). Cuvette temperature was maintained at 16 °C with cooling plates.

## References

- Agatsuma, Y. & Lawrence, J.M. (2013). Chapter 35 - *Hemicentrotus pulcherrimus*, *Pseudocentrotus depressus*, and *Heliocidaris crassispina*. In: *Sea Urchins: Biology and Ecology*. Elsevier, pp. 643-659.
- Aldred, N. & Clare, A.S. (2014). Mini-review: impact and dynamics of surface fouling by solitary and compound ascidians. *Biofouling*, 30, 259-270.
- Almeida, Â., Freitas, R., Calisto, V., Esteves, V.I., Schneider, R.J., Soares, A.M.V.M. *et al.* (2018). Effects of carbamazepine and cetirizine under an ocean acidification scenario on the biochemical and transcriptome responses of the clam *Ruditapes philippinarum*. *Environ Pollut*, 235, 857-868.
- Aluigi, M.G., Diaspro, A., Ramoino, P., Russo, P. & Falugi, C. (2012). The sea urchin, *Paracentrotus lividus*, as a model to investigate the onset of molecules immunologically related to the  $\alpha$ -7 subunit of nicotinic receptors during embryonic and larval development. *Curr Drug Targets*, 13, 587-593.
- Aluigi, M.G., Falugi, C., Mugno, M.G., Privitera, D. & Chiantore, M. (2010). Dose-dependent effects of chlorpyrifos, an organophosphate pesticide, on metamorphosis of the sea urchin, *Paracentrotus lividus*. *Ecotoxicology*, 19, 520-529.
- Backhaus, T. (2014). Medicines, shaken and stirred: a critical review on the ecotoxicology of pharmaceutical mixtures. *Phil Trans R Soc Lond B Biol Sci*, 369, 20130585.
- Barbosa, D.B., Mello, A.d.A., Allodi, S. & de Barros, C.M. (2018). Acute exposure to water-soluble fractions of marine diesel oil: Evaluation of apoptosis and oxidative stress in an ascidian. *Chemosphere*, 211, 308-315.
- Bednaršek, N., Tarling, G.A., Bakker, D.C.E., Fielding, S., Jones, E.M., Venables, H.J. *et al.* (2012). Extensive dissolution of live pteropods in the Southern Ocean. *Nat Geosci*, 5, 881-885.
- Bennett, C.E. & Marshall, D.J. (2005). The relative energetic costs of the larval period, larval swimming and metamorphosis for the ascidian *Diplosoma listerianum*. *Mar Freshw Behav Physiol*, 38, 21-29.
- Berrill, N.J. & Watson, D.M.S. (1935). VIII - Studies in Tunicate development - Part III - Differential retardation and acceleration. *Phil Trans R Soc Lond B Biol Sci*, 225, 255-326.
- Bostwick, M., Smith, E.L., Borba, C., Newman-Smith, E., Guleria, I., Kourakis, M.J. *et al.* (2020). Antagonistic inhibitory circuits integrate visual and gravitactic behaviors. *Curr Biol*, 30, 600-609.e602.
- Boyd, P.W., Collins, S., Dupont, S., Fabricius, K., Gattuso, J.-P., Havenhand, J. *et al.* (2018). Experimental strategies to assess the biological ramifications of multiple drivers of global ocean change—A review. *Glob Change Biol*, 24, 2239-2261.
- Brooks, B.W., Turner, P.K., Stanley, J.K., Weston, J.J., Glidewell, E.A., Foran, C.M. *et al.* (2003). Waterborne and sediment toxicity of fluoxetine to select organisms. *Chemosphere*, 52, 135-142.
- Brown, E.R., Nishino, A., Bone, Q., Meinertzhagen, I.A. & Okamura, Y. (2005). GABAergic synaptic transmission modulates swimming in the ascidian larva. *Eur J Neurosci*, 22, 2541-2548.

- Bryant, V., Newbery, D.M., McLusky, D.S. & Campbell, R. (1985). Effect of temperature and salinity on the toxicity of nickel and zinc to two estuarine invertebrates (*Corophium volutator*, *Macoma balthica*). *Mar Ecol Prog Ser*, 24, 139-153.
- Burke, R.D., Angerer, L.M., Elphick, M.R., Humphrey, G.W., Yaguchi, S., Kiyama, T. *et al.* (2006). A genomic view of the sea urchin nervous system. *Dev Biol*, 300, 434-460.
- Buznikov, G.A., Lambert, H.W. & Lauder, J.M. (2001). Serotonin and serotonin-like substances as regulators of early embryogenesis and morphogenesis. *Cell Tissue Res*, 305, 177-186.
- Byrne, M. & Przeslawski, R. (2013). Multistressor Impacts of Warming and Acidification of the Ocean on Marine Invertebrates' Life Histories. *Integr Compar Biol*, 53, 582-596.
- Caldeira, K. & Wickett, M.E. (2005). Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean. *J Geophys Res*, 110.
- Chan, K.Y., Grünbaum, D. & O'Donnell, M.J. (2011). Effects of ocean-acidification-induced morphological changes on larval swimming and feeding. *J Exp Biol*, 214, 3857-3867.
- Chan, K.Y.K., García, E. & Dupont, S. (2015a). Acidification reduced growth rate but not swimming speed of larval sea urchins. *Sci Rep*, 5, 9764.
- Chan, K.Y.K. & Grünbaum, D. (2010). Temperature and diet modified swimming behaviors of larval sand dollar. *Mar Ecol Prog Ser*, 415, 49-59.
- Chan, K.Y.K., Grünbaum, D., Arnberg, M. & Dupont, S. (2015b). Impacts of ocean acidification on survival, growth, and swimming behaviours differ between larval urchins and brittlestars. *ICES J Mar Sci*, 73, 951-961.
- Chan, K.Y.K., Sewell, M.A. & Byrne, M. (2018). Revisiting the larval dispersal black box in the Anthropocene. *ICES Journal of Marine Science*, 75, 1841-1848.
- Chan, K.Y.K. & Tong, C.S.D. (2020). Temporal variability modulates pH impact on larval sea urchin development. *Conserv Physiol*, 8.
- Chiu, S.T. (1990). Age and growth of *Anthocardaris crassispina* (Echinodermata: Echinoida) in Hong Kong. *Bull Mar Sci*, 47, 94-103.
- Chu, S. & Metcalfe, C.D. (2007). Analysis of paroxetine, fluoxetine and norfluoxetine in fish tissues using pressurized liquid extraction, mixed mode solid phase extraction cleanup and liquid chromatography–tandem mass spectrometry. *J Chromatogr A*, 1163, 112-118.
- Corbo, J.C., Di Gregorio, A. & Levine, M. (2001). The Ascidian as a model organism in developmental and evolutionary biology. *Cell*, 106, 535-538.
- Cowen, R.K. & Sponaugle, S. (2009). Larval dispersal and marine population connectivity. *Ann Rev Mar Sci*, 1, 443-466.
- Dahlberg, C., Auger, H., Dupont, S., Sasakura, Y., Thorndyke, M. & Joly, J.-S. (2009). Refining the *Ciona intestinalis* model of central nervous system regeneration. *PLOS ONE*, 4, e4458.
- David, G.K., Marshall, D.J. & Riginos, C. (2010). Latitudinal variability in spatial genetic structure in the invasive ascidian, *Styela plicata*. *Mar Biol*, 157, 1955-1965.



- Deblonde, T., Cossu-Leguille, C. & Hartemann, P. (2011). Emerging pollutants in wastewater: A review of the literature. *The second European PhD students workshop: Water and health, Cannes 2010*, 214, 442-448.
- Dehal, P., Satou, Y., Campbell, R.K., Chapman, J., Degnan, B., De Tomaso, A. *et al.* (2002). The draft genome of *Ciona intestinalis*: Insights into chordate and vertebrate origins. *Science*, 298, 2157-2167.
- Di Poi, C., Darmaillacq, A.-S., Dickel, L., Boulouard, M. & Bellanger, C. (2013). Effects of perinatal exposure to waterborne fluoxetine on memory processing in the cuttlefish *Sepia officinalis*. *Aquat Toxicol*, 132-133, 84-91.
- Di Poi, C., Evariste, L., Serpentine, A., Halm-Lemeille, M.P., Lebel, J.M. & Costil, K. (2014). Toxicity of five antidepressant drugs on embryo–larval development and metamorphosis success in the Pacific oyster, *Crassostrea gigas*. *Environ Sci Pollut Res Int*, 21, 13302-13314.
- Dickson, A.G. & Millero, F.J. (1987). A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res Part I Oceanogr Res Pap*, 34, 1733-1743.
- Dorey, N., Maboloc, E. & Chan, K.Y.K. (2018). Development of the sea urchin *Heliocidaris crassispina* from Hong Kong is robust to ocean acidification and copper contamination. *Aquat Toxicol*, 205, 1-10.
- Eguchi, Y., Shimizu, S. & Tsujimoto, Y. (1997). Intracellular ATP levels determine cell death fate by apoptosis or necrosis. *Cancer Res*, 57, 1835-1840.
- Elmore, S. (2007). Apoptosis: A review of programmed cell death. *Toxicol Pathol*, 35, 495-516.
- Epel, D., Cole, B., Hamdoun, A. & Thurber, R.V. (2006). The sea urchin embryo as a model for studying efflux transporters: Roles and energy cost. *Pollutant Responses in Marine Organisms (PRIMO 13)*, 62, S1-S4.
- Fabbri, E. & Franzellitti, S. (2016). Human pharmaceuticals in the marine environment: Focus on exposure and biological effects in animal species. *Environ Toxicol Chem*, 35, 799-812.
- Fenet, H., Arpin-Pont, L., Vanhoutte-Brunier, A., Munaron, D., Fiandrino, A., Martínez Bueno, M.-J. *et al.* (2014). Reducing PEC uncertainty in coastal zones: A case study on carbamazepine, oxcarbazepine and their metabolites. *Environ Int*, 68, 177-184.
- Fent, K., Weston, A.A. & Caminada, D. (2006). Ecotoxicology of human pharmaceuticals. *Aquat Toxicol*, 76, 122-159.
- Fong, P.P., Bury, T.B., Dworkin-Brodsky, A.D., Jasion, C.M. & Kell, R.C. (2015). The antidepressants venlafaxine (“Effexor”) and fluoxetine (“Prozac”) produce different effects on locomotion in two species of marine snail, the oyster drill (*Urosalpinx cinerea*) and the starsnail (*Lithopoma americanum*). *Mar Environ Res*, 103, 89-94.
- Ford, A.T., Hyett, B., Cassidy, D. & Malyon, G. (2018). The effects of fluoxetine on attachment and righting behaviours in marine (*Gibbula unibilicalis*) and freshwater (*Lymnaea stagnalis*) gastropods. *Ecotoxicology*, 27, 477-484.
- Ford, A.T., Ågerstrand, M., Brooks, B.W., Allen, J., Bertram, M.G., Brodin, T. *et al.* (2021). The role of behavioral ecotoxicology in environmental protection. *Environ Sci Technol*, 55, 5620-5628.

- Freitas, R., Almeida, Â., Calisto, V., Velez, C., Moreira, A., Schneider, R.J. *et al.* (2016). The impacts of pharmaceutical drugs under ocean acidification: New data on single and combined long-term effects of carbamazepine on *Scrobicularia plana*. *Sci Total Environ*, 541, 977-985.
- Fuchs, H.L. & Gerbi, G.P. (2016). Seascape-level variation in turbulence- and wave-generated hydrodynamic signals experienced by plankton. *Prog Oceanogr*, 141, 109-129.
- Förstner, U. & Wittmann, G.T.W. (1981). *Metal Pollution in the Aquatic Environment*. 2 edn. Springer-Verlag Berlin Heidelberg.
- Gallo, A. & Tosti, E. (2015). The ascidian *Ciona intestinalis* as model organism for ecotoxicological assays. *J Marine Sci Res Dev*, 5, 1000e1138.
- Gaw, S., Thomas, K.V. & Hutchinson, T.H. (2014). Sources, impacts and trends of pharmaceuticals in the marine and coastal environment. *Philos Trans R Soc Lond B Biol Sci*, 369.
- Geissen, V., Mol, H., Klumpp, E., Umlauf, G., Nadal, M., van der Ploeg, M. *et al.* (2015). Emerging pollutants in the environment: A challenge for water resource management. *Int Soil Water Conserv Res*, 3, 57-65.
- Gelatti, U., Pedrazzani, R., Marcantoni, C., Mascaretti, S., Repice, C., Filippucci, L. *et al.* (2013). ‘You’ve got m@il: Fluoxetine coming soon!’: Accessibility and quality of a prescription drug sold on the web. *Int J Drug Policy*, 24, 392-401.
- Gerhardt, A. (2007). Aquatic behavioral ecotoxicology-prospects and limitations. *Hum Ecol Risk Assess*, 13, 481-491.
- Grünbaum, D. & Strathmann, R.R. (2003). Form, performance and trade-offs in swimming and stability of armed larvae. *J Mar Res*, 61, 659-691.
- Guler, Y. & Ford, A.T. (2010). Anti-depressants make amphipods see the light. *Aquat Toxicol*, 99, 397-404.
- Gulkowska, A., He, Y., So, M.K., Yeung, L.W., Leung, H.W., Giesy, J.P. *et al.* (2007). The occurrence of selected antibiotics in Hong Kong coastal waters. *Mar Pollut Bull*, 54, 1287-1293.
- Hamilton, T.J., Kwan, G.T., Gallup, J. & Tresguerres, M. (2016). Acute fluoxetine exposure alters crab anxiety-like behaviour, but not aggressiveness. *Sci Rep*, 6, 19850.
- Hen, R. (1993). Structural and functional conservation of serotonin receptors throughout evolution. *Exs*, 63, 266-278.
- Hird, C.M., Urbina, M.A., Lewis, C.N., Snape, J.R. & Galloway, T.S. (2016). Fluoxetine exhibits pharmacological effects and trait-based sensitivity in a marine worm. *Environ Sci Technol*, 50, 8344-8352.
- Hofmann, G.E., Barry, J.P., Edmunds, P.J., Gates, R.D., Hutchins, D.A., Klinger, T. *et al.* (2010). The effect of ocean acidification on calcifying organisms in marine ecosystems: An organism-to-ecosystem perspective. *Ann Rev Ecol Evol Syst*, 41, 127-147.
- Hofmann, G.E., Smith, J.E., Johnson, K.S., Send, U., Levin, L.A., Micheli, F. *et al.* (2011). High-frequency dynamics of ocean pH: a multi-ecosystem comparison. *PLoS One*, 6, e28983.
- Horie, T., Nakagawa, M., Sasakura, Y. & Kusakabe, T.G. (2009). Cell type and function of neurons in the ascidian nervous system. *Dev Growth Differ*, 51, 207-220.

- Katow, H., Abe, K., Katow, T., Zamani, A. & Abe, H. (2013). Development of the GABA-ergic signaling system and its role in larval swimming in sea urchin. *J Exp Biol*, 216, 1704-1716.
- Kolpin, D.W., Furlong, E.T., Meyer, M.T., Thurman, E.M., Zaugg, S.D., Barber, L.B. *et al.* (2002). Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams, 1999–2000: A National Reconnaissance. *Environ Sci Technol*, 36, 1202-1211.
- Kot-Wasik, A., Jakimska, A. & Śliwka-Kaszyńska, M. (2016). Occurrence and seasonal variations of 25 pharmaceutical residues in wastewater and drinking water treatment plants. 188, 661.
- Kwon, J.-W. & Armbrust, K.L. (2006). Laboratory persistence and fate of fluoxetine in aquatic environments. *Environ Toxicol Chem*, 25, 2561-2568.
- Lazzara, R., Blázquez, M., Porte, C. & Barata, C. (2012). Low environmental levels of fluoxetine induce spawning and changes in endogenous estradiol levels in the zebra mussel *Dreissena polymorpha*. 106-107, 123-130.
- Lemaire, P., Smith, W.C. & Nishida, H. (2008). Ascidians and the Plasticity of the Chordate Developmental Program. *Curr Biol*, 18, R620-R631.
- Lo, H.K.A. (2020). Impacts of emerging pollutants on early development of marine invertebrates: case studies on microplastics and interaction between fluoxetine and ocean acidification. The Hong Kong University of Science and Technology.
- Lo, H.K.A., Chua, V.A. & Chan, K.Y.K. (in review). Near future ocean acidification modulates the impact of fluoxetine at environmental concentration on larval urchins. *Sci Total Environ*.
- Lynch, T. & Price, A. (2007). The effect of cytochrome P450 metabolism on drug response, interactions, and adverse effects. *Am Fam Physician*, 76, 391-396.
- Maboloc, E.A., Batzel, G., Grünbaum, D. & Chan, K.Y.K. (2020). Vertical distribution of echinoid larvae in pH stratified water columns. *Mar Biol*, 167, 13.
- McHenry, M.J. (2005). The morphology, behavior, and biomechanics of swimming in ascidian larvae. *Can J Zool*, 83, 62-74.
- Mehrbach, C., Culbertson, C.H., Hawley, J.E. & Pytkowicz, R.M. (1973). Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure. *Limnol Oceanogr*, 18, 897-907.
- Minchinton, T.E. & Scheibling, R.E. (1991). The influence of larval supply and settlement on the population structure of barnacles. *Ecology*, 72, 1867-1879.
- Mogami, Y., Watanabe, K., Ooshima, C., Kawano, A. & Baba, S.A. (1992). Regulation of ciliary movement in sea urchin embryos: Dopamine and 5-HT change the swimming behaviour. *Comp Biochem Physiol C Toxicol*, 101, 251-254.
- Mojtabai, R. & Olfson, M. (2014). National trends in long-term use of antidepressant medications: results from the US National Health and Nutrition Examination Survey. *J Clin Psychiatry*, 75, 169-177.
- Monteiro, S.C. & Boxall, A.B.A. (2010). Occurrence and fate of human pharmaceuticals in the environment. In: *Reviews of Environmental Contamination and Toxicology*. Springer New York New York, NY, pp. 53-154.
- Méndez, N. & Barata, C. (2015). Effects of the antidepressant fluoxetine in spiked-sediments on developmental and reproductive features of the polychaetes *Capitella teleta* and *Capitella* sp A. *Ecotoxicology*, 24, 106-118.

- Nakagawa, M., Miyamoto, T., Ohkuma, M. & Tsuda, M. (1999). Action spectrum for the photophobic response of *Ciona intestinalis* (Ascidiacea, Urochordata) larvae implicates retinal protein. *Photochem Photobiol*, 70, 359-362.
- Nakamura, Y., Yamamoto, H., Sekizawa, J., Kondo, T., Hirai, N. & Tatarazako, N. (2008). The effects of pH on fluoxetine in Japanese medaka (*Oryzias latipes*): Acute toxicity in fish larvae and bioaccumulation in juvenile fish. *Chemosphere*, 70, 865-873.
- Naranjo, S.A., Carballo, J.L. & García-Gómez, J.C. (1996). Effects of environmental stress on ascidian populations in Algeciras Bay (southern Spain). Possible marine bioindicators? *Mar Ecol Prog Ser*, 144, 119-131.
- Nilsson, G.E., Dixon, D.L., Domenici, P., McCormick, M.I., Sørensen, C., Watson, S.-A. *et al.* (2012). Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nat Clim Change*, 2, 201-204.
- Okamura, Y., Nishino, A., Murata, Y., Nakajo, K., Iwasaki, H., Ohtsuka, Y. *et al.* (2005). Comprehensive analysis of the ascidian genome reveals novel insights into the molecular evolution of ion channel genes. *Physiol Genomics*, 22, 269-282.
- Pearse, J.S. (2006). Ecological role of purple sea urchins. *Science*, 314, 940-941.
- Pechenik, J.A. (1999). On the advantages and disadvantages of larval stages in benthic marine invertebrate life cycles. *Mar Ecol Prog Ser*, 177, 269-297.
- Pecquet, A., Dorey, N. & Chan, K.Y.K. (2017). Ocean acidification increases larval swimming speed and has limited effects on spawning and settlement of a robust fouling bryozoan, *Bugula neritina*. *Special Issue: Hong Kong Conference 2016*, 124, 903-910.
- Peters, J.R. & Granek, E.F. (2016). Long-term exposure to fluoxetine reduces growth and reproductive potential in the dominant rocky intertidal mussel, *Mytilus californianus*. *Sci Total Environ*, 545-546, 621-628.
- Peters, J.R., Granek, E.F., de Rivera, C.E. & Rollins, M. (2017). Prozac in the water: Chronic fluoxetine exposure and predation risk interact to shape behaviors in an estuarine crab. *Ecol Evol*, 7, 9151-9161.
- Pineda, M.C., López-Legentil, S. & Turon, X. (2011). The whereabouts of an ancient wanderer: global phylogeography of the solitary ascidian *Styela plicata*. *PLOS ONE*, 6, e25495.
- Polverino, G., Martin, J.M., Bertram, M.G., Soman, V.R., Tan, H., Brand, J.A. *et al.* (2021). Psychoactive pollution suppresses individual differences in fish behaviour. *Proc R Soc B Biol Sci*, 288, 20202294.
- Rand-Weaver, M., Margiotta-Casaluci, L., Patel, A., Panter, G.H., Owen, S.F. & Sumpter, J.P. (2013). The read-across hypothesis and environmental risk assessment of pharmaceuticals. *Environ Sci Technol*, 47, 11384-11395.
- Razy-Krajka, F., Brown, E.R., Horie, T., Callebert, J., Sasakura, Y., Joly, J.-S. *et al.* (2012). Monoaminergic modulation of photoreception in ascidian: evidence for a proto-hypothalamo-retinal territory. *BMC Biol*, 10, 45.
- Ricke, K.L. & Caldeira, K. (2014). Maximum warming occurs about one decade after a carbon dioxide emission. *Environ Res Lett*, 9, 124002.
- Roy, A., Metaxas, A. & Ross, T. (2012). Swimming patterns of larval *Strongylocentrotus droebachiensis* in turbulence in the laboratory. *Mar Ecol Prog Ser*, 453, 117-127.

- Rudolf, J., Dondorp, D., Canon, L., Tiew, S. & Chatzigeorgiou, M. (2019). Automated behavioural analysis reveals the basic behavioural repertoire of the urochordate *Ciona intestinalis*. *Sci Rep*, 9, 2416.
- Sarà, G., Mangano, M.C., Johnson, M. & Mazzola, A. (2018). Integrating multiple stressors in aquaculture to build the blue growth in a changing sea. *Hydrobiologia*, 809, 5-17.
- Satoh, N. (1994). *Developmental Biology of Ascidiaceans*. Cambridge University Press, Cambridge.
- Singh, S.P., Azua, A., Chaudhary, A., Khan, S., Willett, K.L. & Gardinali, P.R. (2010). Occurrence and distribution of steroids, hormones and selected pharmaceuticals in South Florida coastal environments. *Ecotoxicology*, 19, 338-350.
- Smith, E.M., Iftikar, F.I., Higgins, S., Irshad, A., Jandoc, R., Lee, M. *et al.* (2012). In vitro inhibition of cytochrome P450-mediated reactions by gemfibrozil, erythromycin, ciprofloxacin and fluoxetine in fish liver microsomes. *Aquat Toxicol*, 109, 259-266.
- Sokolova, I.M., Frederich, M., Bagwe, R., Lannig, G. & Sukhotin, A.A. (2012). Energy homeostasis as an integrative tool for assessing limits of environmental stress tolerance in aquatic invertebrates. *Mar Environ Res*, 79, 1-15.
- Song, F., Freemantle, N., Sheldon, T.A., House, A., Watson, P., Long, A. *et al.* (1993). Selective serotonin reuptake inhibitors: meta-analysis of efficacy and acceptability. *Bri Med J*, 306, 683-687.
- Stockdale, A., Tipping, E., Lofts, S. & Mortimer, R.J.G. (2016). Effect of ocean acidification on organic and inorganic speciation of trace metals. *Environ Sci Technol*, 50, 1906-1913.
- Strathmann, M.F. (1987). *Reproduction and Development of Marine Invertebrates of the Northern Pacific Coast: Data and Methods for the Study of Eggs, Embryos, and Larvae*. University of Washington Press.
- Strathmann, R.R. & Grünbaum, D. (2006). Good eaters, poor swimmers: compromises in larval form. *Integr Compar Biol*, 46, 312-322.
- Stumpff, M., Hu, M.Y., Melzner, F., Gutowska, M.A., Dorey, N., Himmerkus, N. *et al.* (2012). Acidified seawater impacts sea urchin larvae pH regulatory systems relevant for calcification. *Proc Natl Acad Sci U.S.A.*, 109, 18192-18197.
- Todgham, A.E. & Hofmann, G.E. (2009). Transcriptomic response of sea urchin larvae *Strongylocentrotus purpuratus* to CO<sub>2</sub>-driven seawater acidification. *J Expl Biol*, 212, 2579-2594.
- True, A.C., Webster, D.R., Weissburg, M.J., Yen, J. & Genin, A. (2015). Patchiness and depth-keeping of copepods in response to simulated frontal flows. *Mar Ecol Prog Ser*, 539, 65-76.
- Tsuda, M., Kawakami, I. & Shiraishi, S. (2003). Sensitization and habituation of the swimming behavior in ascidian larvae to light. *Zool Sci*, 20, 13-22, 10.
- Tunnickliff, G., Schindler, N.L., Crites, G.J., Goldenberg, R., Yochum, A. & Malatynska, E. (1999). The GABA(A) receptor complex as a target for fluoxetine action. *Neurochem Res*, 24, 1271-1276.
- van Grinsven, H.J.M., ten Berge, H.F.M., Dalgaard, T., Fraters, B., Durand, P., Hart, A. *et al.* (2012). Management, regulation and environmental impacts of nitrogen

- fertilization in northwestern Europe under the Nitrates Directive; a benchmark study. *Biogeosciences*, 9, 5143-5160.
- Wada, Y., Mogami, Y. & Baba, S. (1997). Modification of ciliary beating in sea urchin larvae induced by neurotransmitters: beat-plane rotation and control of frequency fluctuation. *J Exp Biol*, 200, 9-18.
- Wang, L., Li, Q., Bi, H. & Mao, X.Z. (2016). Human impacts and changes in the coastal waters of south China. *Sci Total Environ*, 562, 108-114.
- Webb, S.F. (2004). A Data-based Perspective on the Environmental Risk Assessment of Human Pharmaceuticals I — Collation of Available Ecotoxicity Data. In: *Pharmaceuticals in the Environment: Sources, Fate, Effects and Risks* (ed. Kümmerer, K). Springer Berlin Heidelberg Berlin, Heidelberg, pp. 317-343.
- Weston, J.J., Huggett, D., Rimoldi, J., Foran, C.M. & Statter, M. (2001). Determination of fluoxetine (Prozac™) and norfluoxetine in the aquatic environment. Annual Meeting of the Society of Environmental Toxicology and Chemistry, Baltimore, MD. 142, 169-183.
- Wobbrock, J.O., Findlater, L., Gergle, D. & Higgins, J.J. (2011). The aligned rank transform for nonparametric factorial analyses using only ANOVA procedures. In: *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems*. Association for Computing Machinery Vancouver, BC, Canada, pp. 143–146.
- Yamamoto, H., Hayashi, A., Nakamura, Y. & Sekizawa, J. (2005). Fate and partitioning of selected pharmaceuticals in aquatic environment. *Environ Sci*, 12, 347-358.
- Zega, G., Thorndyke, M.C. & Brown, E.R. (2006). Development of swimming behaviour in the larva of the ascidian *Ciona intestinalis*. *J Exp Biol*, 209, 3405-3412.
- Zindler, F., Tisler, S., Loerracher, A.-K., Zwiener, C. & Braunbeck, T. (2020). Norfluoxetine is the only metabolite of fluoxetine in zebrafish (*Danio rerio*) embryos that accumulates at environmentally relevant exposure scenarios. *Environ Sci Technol*, 54, 4200-4209.